Circulating Permeability Factors in Idiopathic Nephrotic Syndrome

Tae-Sun Ha, M.D.

Department of Pediatrics, College of Medicine and Medical Research Institute, Chungbuk National University, Cheongju, Korea

Corresponding author: Tae-Sun Ha, M.D.
Department of Pediatrics, College of Medicine, Chungbuk National University, Department of Pediatrics, Chungbuk National University Hospital, 1 Chungdae-ro, Seowon-gu, Cheongju, Korea 28644,
Tel: +82-43-269-6374
Fax: +82-43-264-6620
E-mail: tsha@chungbuk.ac.kr

Received: 26 March 2019
Revised: 29 March 2019
Accepted: 11 April 2019

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2019 The Korean Society of Pediatric Nephrology

Nephrotic syndrome (NS) is a common chronic glomerular disease in children characterized by significant proteinuria with resulting hypoalbuminemia, edema, and hyperlipidemia. Renal biopsy findings of diffuse foot processes effacement on electron microscopy and minimal change disease, focal segmental glomerulosclerosis (FSGS), or diffuse mesangial proliferation on light microscopy. It has been speculated that circulating permeability factors would be implicated in the pathogenesis of NS because they have been reportedly detected in the sera of patients and in experimental models of induced proteinuria. Moreover, a substantial portion of the patients with primary FSGS recurrence shortly after transplantation. This report reviews the current knowledge regarding the role of circulating permeability factors in the pathogenesis of proteinuria in NS and suggests future targeted therapeutic approaches for NS.

Key words: Nephrotic syndrome, Permeability Factors, Proteinuria, Minimal change disease, Focal segmental glomerulosclerosis

Introduction

Nephrotic syndrome (NS) is characterized by significant proteinuria (>40 mg/m²/h or a spot urinary protein-to-creatinine ratio of over 2 mg/mg) and hypoalbuminemia (<3.0 g/dL), which in turn causes edema and hyperlipidemia¹⁻⁴. The most common form (90%) of NS in children is primary (idiopathic) and is classified, on the basis of clinical response to steroid therapy, as steroid sensitive or steroid resistant. In terms of pathologic findings, the effacement of diffuse foot processes is common on electron microscopy. Over 80% of childhood patients with NS show minimal change disease (MCD) characterized by normal renal histology on light microscopy, where the remaining 20% show focal segmental glomerulosclerosis (FSGS), diffuse mesangial proliferation, or mesangial proliferative glomerulonephritis¹⁻⁴.

Recent advances in podocyte biology have enforced our knowledge of the molecular changes and genetic defects of the glomerular filtration structures, which could explain the mechanism of proteinuria in NS⁵⁻⁷. The development of proteinuria in NS is caused by an increased glomerular permeability to plasma proteins, which is accompanied with phenotypical changes of podocytes. The Shalhoub hypothesis⁸ suggests that circulating permeability factors could be implicated in the structural and molecular changes of podocytes.

cytes in idiopathic NS. However, defining a single putative factor remains elusive¹⁻⁴. This paper reviews the current experimental and clinical knowledge of the role of circulating permeability factors in the pathogenesis of proteinuria in idiopathic NS.

Evidence of circulating permeability factors in NS

The pathologic role of circulating permeability factors in podocyte structure and function, leading to proteinuria and NS, is supported by the following clinical and experimental observations^{9,10}: (1) a child born from a mother with FSGS and NS spontaneously recovered from proteinuria after maternal transfer of nephrotic proteinuria^{11,12}; (2) the perfusion of isolated rat glomeruli with plasma of patients with FSGS induced an increased glomerular permeability to albumin as compared to normal control plasma^{13,14}; (3) after kidney transplantation from a donor without the disease, approximately 30% of patients with FSGS developed massive proteinuria within hours to days, followed by FSGS histological lesions¹⁵⁻¹⁷; (4) patients with recurrent FSGS after kidney transplantation showed complete or partial response to early plasma exchange¹⁸⁻²¹; (5) pre-transplant plasmapheresis or immunotherapy reduced the risk of FSGS recurrence after transplantation^{22,23}; (6) the successful transplantation in an end-stage renal disease patient due to diabetic nephropathy of a graft removed from a transplanted FSGS recipient because of intractable recurrence of massive proteinuria and renal insufficiency²⁴; (7) the kidneys from a donor with FSGS transplanted into two uremic recipients remained free from proteinuria with normal renal function after one year²⁵.

Identification

The clinical and experimental evidence of circulating permeability factors suggest that they are involved in the pathogenesis of glomerular hyperpermeability. Maas et al.²⁶ suggested criteria to define the pathogenic circulating factors in idiopathic MCD and FSGS. In recent decades, research has focused on identifying permeability factors

using various models, as will be discussed in the following sections.

Isolation of responsible factors

In 1974, Shalhoub⁸ theorized that a disordered clone of T-lymphocytes, present in both MCD and FSGS, secreted a circulating lymphokine “toxic” to the glomerular barrier. The first experimental evidence suggesting the pathophysiological role of lymphokine in idiopathic NS was reported by Lagrue et al.²⁷ in 1975, showing that the supernatant of lymphocytes from patients with MCD elaborated *in vitro* by concanavalin A-stimulated T-lymphocytes contained a factor that modified the vascular permeability. The molecular characteristics of the permeability factors have been derived from findings that the active fraction of sera from patients with FSGS precipitates in 70-80% ammonium sulfate solution independent of the immunoglobulin fraction. The putative permeability factors were found to bind to protein A and had a molecular size between 30 and 50 kDa²⁸.

In vitro experiments

The role of isolated permeability factors could be examined by *in vitro* endothelial cell or podocyte culture models. Observations of cultured found changes in the configuration of the actin cytoskeleton, cell morphology, and protein expression in responses to permeability factors. Sera from patients with FSGS recurrence were found to disrupt podocyte focal complexes via immunofluorescence²⁹. Hemopexin was found to affect the podocyte actin cytoskeleton and reduced endothelial glycocalyx, thereby increasing albumin diffusion^{30,31}.

Ex vivo experiments

Glomerular permeability could be measured using *ex vivo* isolated glomeruli developed by Savin et al.³². In isolated rat glomeruli, when an isotonic albumin oncotic solution was replaced by a solution with a lower albumin concentration, an increase in glomerular size was observed

through swelling when the permeability barrier was intact. It was found that serum from patients with FSGS increased the permeability of glomeruli to albumin²⁸⁾. Recently, Desideri et al.³³⁾ improved the *ex vivo* model and found that incubation with plasma from patients with post-transplant recurrence of NS increased albumin permeability in rat glomeruli compared to remission plasma. The blocking effect of galactose on circulating permeability factors was demonstrated using *ex vivo* isolated glomeruli³⁴⁾. A glomerulus-on-a-chip microdevice is a recently developed *ex vivo* model, which could replace isolated glomeruli³⁵⁾.

In vivo experiments

Plasma permeability factors have been found to induce proteinuria in experimental animals. Zimmermann¹³⁾ demonstrated that serum from patients with recurrent FSGS after renal transplantation could induce increased albumin

excretion when infused in the aorta of intact rats. Koyama et al.³⁶⁾ found that human T cell hybridomas formed from cells from patients with MCD produced a substance that induced proteinuria and reduced the density of polyethylene staining of glomerular anionic sites when injected intravenously into normal Wistar rats. Each permeability factor, such as hemopexin³⁷⁾ and cardiotrophin-like cytokine factor-1 (CLCF-1)³⁸⁾, also induced proteinuria when injected intravenously into normal rats.

Suggested permeability factors

Although the clinical and experimental data suggest that a circulating permeability factor induces glomerular proteinuria and NS, a single putative factor has yet to be identified. Several candidates are implicated in the development of proteinuria in idiopathic MCD and FSGS^{9,10,26,39-41)}. The proposed candidates are listed in Table 1. Clinical and ex-

Table 1. Circulating Permeability Factors Proposed in Idiopathic Nephrotic Syndrome

Circulating factor	Molecular weight (kDa)	Major targets	Preferences
VPF	60–160	Anionic charges Endothelium	MCD FSGS recurrence FSGS
FSGS factor	30–50	Podocyte	FSGS FSGS recurrence
Hemopexin	54–70	Anionic charges Podocyte Endothelium	MCD
CD80 (B7-1)	55–60	Podocyte	MCD FSGS recurrence FSGS
suPAR	20–50	Podocyte Integrin	FSGS FSGS recurrence
CLCF-1	22	Podocyte	FSGS FSGS recurrence
Angptl-4	45–65	Anionic charges	MCD FSGS
IL-13	10–17	Podocyte Adaptors	MCD
IL-8	11	Anionic charges	MCD
CD40 auto-antibody	150	Podocyte Integrin	FSGS recurrence
Heparanase	57–60	Anionic charges	MCD FSGS

Abbreviations: VPF, vascular permeability factor; FSGS, focal segmental glomerular sclerosis; CD, Cluster of Differentiation; suPAR, soluble urokinase plasminogen activator receptor; CLCF-1, cardiotrophin-like cytokine factor-1; angptl-4, angiopoietin like-4; IL-13, Interleukin-13; IL-8, Interleukin-8; MCD, minimal change disease.

perimental data have suggested roles for cytokine interleukin (IL)-13, CD80, anti-CD40 antibody, angiopoietin-like 4 (angptl-4), circulating CLCF-1 (member of the IL-6 family), circulating hemopexin, and soluble urokinase-type plasminogen activator receptor (suPAR), among others, in the development of NS.

Vascular permeability factor (VPF)

Lagrué et al.²⁷⁾ were the first to systematically demonstrate the existence of a factor that could modulate vascular permeability. They defined potential plasma factors on vascular permeability after finding that VPF was produced *in vitro* after the stimulation of isolated blood T-lymphocytes with concanavalin A. Lymphocyte-derived VPF following concanavalin A stimulation *in vitro* was found to reduce glomerular polyanions by colloidal iron staining following *in vitro* incubation of kidney cryostat sections, and was found to cause proteinuria in rats^{42,43)}. Concanavalin A-derived supernatants of lymphocytes from patients with MCD induced proteinuria in rats *in vivo* following infusion into the renal artery⁴⁴⁾ or tail vein^{36,45)}. However, lymphocyte-derived VPF has failed to be characterized despite the continuous production of supernatants. Although VPF was first identified as a permeability factor in MCD, the following studies actually showed that VPF was present in most but not all patients with MCD and was not unique to MCD. Moreover, there is no direct evidence that VPF causes proteinuria or that it even has an effect on glomerular permeability²⁶⁾.

FSGS factor

The FSGS factor was identified from observations wherein the active fraction of sera from patients with recurrent FSGS was found to induce proteinuria in rats¹³⁾. Sera from patients with recurrent FSGS increased glomerular permeability to albumin compared to normal control plasma in the perfusion of isolated rat glomeruli model^{14,28)}, and was found to disrupt podocyte focal complexes²⁹⁾. The active fraction of permeability factor precipitates in 70–80% ammonium sulfate solution, inconsistent with immunoglo-

bulin fraction. This putative FSGS factor has a molecular size between 30 and 50 kDa and is bound to protein A²⁸⁾, which is consistent with a report of proteinuria remission after protein A column *ex vivo* adsorption²⁰⁾. This FSGS factor was partly identified as CLCF-1 by sequential purification and fractionation.

Hemopexin

Hemopexin is a serine protease predominantly produced in the liver. It has been suggested that, in normal conditions, circulating hemopexin is inactive; however, under certain circumstances, hemopexin becomes activated as a serine protease⁴⁶⁾. Activated hemopexin appears to alter the function of both glomerular endothelium and podocytes. *In vivo*, activated hemopexin injection in rats caused reversible proteinuria with podocyte foot processes fusion^{37,46)}. In cultured podocytes, hemopexin activated protein kinase B and Ras homolog gene family, member A (RhoA), and induced nephrin-dependent reorganization of the actin cytoskeleton from stress fibers to cytoplasmic aggregates and membrane ruffles^{30,44)}. In addition, hemopexin reduced the anionic sites in the lamina rara interna and endothelial sialoglycoproteins, thereby increasing albumin diffusion across the monolayers of the glomerular endothelial cells^{31,47)}. Studies on human MCD have demonstrated a decreased titer of plasma and urine hemopexin and an increased activity of circulating isoform during disease relapse which then disappeared during remission⁴⁶⁾. These changes were ascribed to an altered configuration of hemopexin in relapsing MCD. No conclusions have been drawn on the role of hemopexin in MCD, and its implication is, at the moment, believed to be a secondary effect.

CD80 (B7-1)

Cluster of differentiation 80 (CD80, also called B7-1) is a transmembrane molecule present on the surface of B cells and other antigen-presenting cells. When bound to its ligand, CD28, present in T-lymphocytes leads to lymphocyte activation⁴⁸⁾. CD80 expression in dendritic cells is inhibited by cytotoxic T-lymphocyte-associated-protein 4 (CTLA-

4) and IL-10, which are produced by T regulatory (Treg) cells^{49,50}. Under normal conditions, CD80 is only transiently expressed after trigger stimuli, and the resulting proteinuria is minimal due to rapid auto-regulatory response of circulating Treg cells or the podocyte itself, leading to the expression of factors such as CTLA-4 to downregulate the podocyte CD80 response⁵¹. The level of circulating CTLA-4 is low during relapses⁵² and there is a predominance of certain CTLA-4 genotypes in idiopathic NS^{53,54}. Furthermore, CD80 has been proposed to be used for differential diagnosis between MCD and FSGS⁵⁵.

The increased expression of CD80 by trigger stimuli results in actin reorganization, as well as a change in the shape of podocytes, leading to the effacement of the foot processes, and proteinuria⁵⁶. Trigger events for increased CD80 podocyte expression in experimental animals include inflammatory stimuli, such as lipopolysaccharide (LPS) and polyinosinic-polycytidylic acid (Poly:IC), oxidative stress, in the case of PAN and IL-13. LPS-injected mice were found to have an increased podocyte CD80 expression through signaling processes mediated by Toll-like receptors (TLR) and IL-13 with foot processes effacement and proteinuria in the early phase which mimicked MCD^{48,57}. In contrast, such findings were not observed in CD80 knockout mice when injected with LPS. Furthermore, severe combined immunodeficient (SCID) mice still developed proteinuria in response to LPS, thus showing that CD80 expression and proteinuria are not dependent on T-lymphocytes⁵⁸. The expression of podocyte CD80 induced by LPS *in vitro* through binding to TLR3 and TLR4⁵⁸ or TNF α treatment⁵⁹ has also been associated with actin derangement and shape changes.

The *in vitro* incubation of human podocytes with poly-IC, a TLR 3 ligand, increased the podocyte expression of CD80 with the activation of TLR3 and cathepsin L, but decreased the expression of synaptopodin, and resulted in actin reorganization via an NF- κ B-dependent pathway, which was partially blocked by dexamethasone⁴⁸. An *In vivo* poly:IC injection also reduced synaptopodin expression in podocytes with proteinuria and increased glomerular CD80 expression and urinary CD80 in mice⁶⁰.

The CD80 levels in podocytes and urine were elevated in MCD patients during the initial onset compared to MCD in remission. CD80 was also detected in the glomeruli of

the same patients but were not found in high levels in other NS-causing nephropathies, such as FSGS or membranous nephropathy (MN)^{61,62}. These results were confirmed in other populations with MCD⁶³, thus leading to the conclusion that high CD80 (B7-1) expression in the urine and glomeruli is a urinary biomarker to distinguish between MCNS and FSGS, which could be used to monitor the activity of the disease^{55,66}. Other studies documented the upregulation of glomerular CD80 (B7-1) in patients with diabetic nephropathy⁶⁴ and/or with post-transplant FSGS recurrence⁶⁵. Sera from MCD patients in relapse, but not in remission, stimulated CD80 expression in cultured podocytes along with diminished nephrin phosphorylation in comparison with podocytes incubated with sera from MCD patients in remission⁶⁶. Therefore, B7-1 may be a biomarker of podocyte injury.

The upregulation of glomerular CD80 (B7-1) in patients with post-transplant FSGS recurrence and active MCD led to the use of abatacept (CTLA-4-Ig), a co-stimulatory inhibitor of B7-1, as a specific therapeutic agent for patients with post-transplant FSGS recurrence and MCD. However, the administration of abatacept resulted in a transient reduction of urinary CD80 levels and resolution of proteinuria in an MCD patient⁶⁷ and substantially reduced proteinuria in FSGS⁶⁵. The results from other studies were also discordant and did not confirm the original observation⁶⁸. The positive data on B7-1 is being challenged by recent discordant findings obtained in clinical data. As such, further controlled studies on the roles of B7-1 and abatacept are needed in various clinical and experimental glomerular pathological conditions.

Soluble urokinase-type plasminogen activator receptor (suPAR)

Urokinase plasminogen activator receptor (uPAR), a cell membrane glycosylphosphatidylinositol (GPI)-anchored membrane glycoprotein, has a role in the migration of activated T-lymphocytes, monocytes, and neutrophils to sites of inflammation⁶⁹⁻⁷¹. uPAR consists of three domains (DI, DII, and DIII), which are encoded by separate exon sets of the *Plaur* gene, and bind to their ligand, urokinase plasminogen activator (uPA)^{9,71,72}. Through interactions with

transcellular receptors, such as integrins, uPAR mediates cell adhesion to vitronectin and activates cell signaling pathways via a mechanism that includes the lipid-dependent activation of $\alpha\beta 3$ integrin for cell migration, proliferation, and survival⁷³. The activation of uPAR is associated with cancer cell invasion and migration⁷⁴. These findings led to predict the role of uPAR in motile podocytes.

Through the cleavage of uPAR from its GPI-anchor, suPAR is released. Further cleavage between the DI and DII/DIII domains of suPAR generates other cleaved suPAR fragments. As suPAR and uPAR are heavily glycosylated proteins, the molecular weight (MW) of uPAR ranges from 35 to 60 kDa, whereas the MW of suPAR ranges from 20 to 50 kDa, depending on the amount of glycosylation and the size of the cleaved proteins^{9,26,39}. Notably, these MW ranges are similar to those of the circulating permeability factor proposed by Savin³⁹.

The suPAR, expressed by several immune cells and endothelial cells, but also by podocytes, can induce foot processes effacement and FSGS *in vitro* and *in vivo* and has been shown to be increased in the plasma of FSGS patients compared to patients with other glomerular diseases, inclusive of MCD^{55,75}. Podocyte foot processes contain an actin-based cytoskeleton that is bound to the vitronectin molecules of the glomerular basement membrane (GBM) via $\alpha 3\beta 1$ and $\alpha\beta 3$ integrins^{71,76}. The activation of vitronectin-dependent $\alpha\beta 3$ -integrin induces intracellular signaling processes, leading to the development of podocyte foot processes and the modifications of the actin cytoskeleton⁷⁶. Podocyte uPAR binds to integrin and vitronectin. In podocyte cultures and murine models, uPAR was found to cause vitronectin-dependent $\alpha\beta 3$ -integrin activation⁷⁷.

Wei et al.⁷⁵ found that the serum concentrations of suPAR were significantly elevated in patients with primary FSGS. However, patients with MCD, MN, and preeclampsia did not display a significant elevation of suPAR levels. In addition, suPAR levels correlated with the presence but not with the level of proteinuria. The highest concentrations of suPAR were found in patients with recurrent FSGS after transplantation. Elevated levels of suPAR prior to transplantation appears to increase the risk of recurrence of disease in the transplanted kidney, and preliminary evidence suggests that treatment with plasmapheresis may significantly reduce these levels and induce remission⁷⁵. Wei et

al.⁷⁸ also defined 3,000 pg/mL as the cut-off value for a high probability of diagnosing FSGS and found that approximately two-thirds of patients with primary FSGS, and to a lesser degree patients with proteinuric kidney disease, were found above this level⁷⁸. In a FSGS CT (70 adults) and Podonet (94 children) cohort study, the authors reported that suPAR levels were elevated in 84.3% (CT cohort) or 55.3% (Podonet cohort) of patients with primary FSGS, and there was an inverse correlation of suPAR levels with estimated glomerular filtration rate (eGFR)⁷⁸. However, in the two studies, the serum suPAR data were not corrected for eGFR. Li et al.⁷⁹ also confirmed in their cohort (109 primary FSGS, 20 MCD, 22 MN, and 96 healthy controls) that suPAR levels were elevated in about half of their patients with FSGS, and could therefore discriminate between FSGS and other proteinuric kidney diseases. In addition, the suPAR levels indicated the steroid-responsiveness of FSGS⁷⁹. However, serum suPAR levels did not differ between primary and secondary FSGS⁸⁰. The initial findings of suPAR as a potential causal factor in primary FSGS was a major breakthrough, and many researchers have subsequently tested the serum suPAR levels in human adult⁸⁰⁻⁸⁴ and pediatric⁸⁵⁻⁸⁷ cohorts with conflicting results after controlling suPAR values for eGFR.

The urinary suPAR level of patients with primary FSGS were also significantly higher compared to those with MCD, MN, secondary FSGS, or normal subjects⁸⁸, and were positively correlated with 24-hour urinary protein excretion in primary FSGS. During the follow-up, the urinary suPAR level of patients with complete remission decreased significantly. After incubation of human podocytes with urine obtained from patients with primary FSGS at presentation, the AP5 signal was induced and then reduced by a blocking antibody specific to uPAR⁸⁸.

The pathological role of suPAR on the development of FSGS has been demonstrated in both *in vivo* and *in vitro* experimental models. Wei et al.⁷⁵ established three different mouse models and confirmed that suPAR caused proteinuria and FSGS in mice by binding to the podocyte $\beta 3$ integrin. In *Plaur*^{-/-} mice, infusion with over 20 μ g of recombinant mouse suPAR protein induced podocyte deposition of suPAR with an increase in $\beta 3$ integrin activity and proteinuria within 24 hours. Furthermore, wild-type mice with transplanted kidneys from *Plaur*^{-/-} mice modeling en-

ogenous suPAR release developed proteinuria after being challenged with LPS, indicating that circulating suPAR may be able to activate $\beta 3$ integrin independently of uPAR. In the third model, wild-type mice with plasmid transfer (*sPlaur*_{WT}, suPARI-II producing plasmid) into the skin showed a rise of serum suPAR and FSGS-like lesions with proteinuria. However, mice that received a gene transfer for the defective $\beta 3$ -integrin binding suPAR mutant, *sPlaur*_{E134A}, did not become albuminuric, suggesting that binding of suPAR to $\beta 3$ integrin is important in suPAR-induced renal injury. Subsequently, the mouse uPAR isoform 2 was found to induce FSGS with proteinuria using suPAR2 transgenic mice, and $\beta 3$ integrin was found to play a critical role in the development of renal dysfunction in suPAR2-transgenic mice in three different integrin $\beta 3$ deficiency models⁸⁹. Alfano et al.⁹⁰ also showed that high dose (20 μ g) of suPAR (full-length recombinant murine suPAR) induced proteinuria in *Plaur*^{-/-} mice and suPAR decreased nephrin expression in podocytes via the suppression of Wilms tumor-1 (WT-1) transcription factor *In vivo* and *in vitro*. Interestingly, only the full-length suPAR molecule interacted with $\beta 3$ -integrin and caused podocyte damage, however, cleaved suPAR molecules were not able to activate $\beta 3$ -integrin. On the other hand, Delville et al.⁹¹ revealed that injecting suPAR (recombinant human suPAR) did not induce proteinuria in wild-type mice unless injected in combination with other podocyte-damaging agents, such as anti-CD40 antibodies. In contrast, Spinale et al.⁸³ did not detect proteinuria in wild-type mice 24 hours after the administration of recombinant suPAR (recombinant mouse suPAR-Fc), despite high suPAR levels. Furthermore, the ectopic expression of the full-length suPAR (*DI-DIII*) molecule from the liver did not induce proteinuria for 44 days despite the elevated suPAR levels. Similarly, Cathelin et al.⁹² found that neither the short-term nor prolonged administration of suPAR (recombinant mouse suPAR-Fc and monomeric mouse uPAR produced in S2-cells) could induce proteinuria in wild-type mice⁵⁹.

Many questions remain regarding the potential pathogenic role of plasma and serum suPAR in primary FSGS. However, from the evidence presented above, suPAR appears to be a biomarker for reduced renal function. Further studies will be needed to clarify the role of suPAR in podocyte damage in primary FSGS patients to provide the

novel treatment strategies for FSGS patients^{9,40,71,93-95}.

Cardiotrophin-like cytokine factor-1 (CLCF-1)

CLCF-1, a member of the IL-6 family of cytokines with a predicted MW of 22 kD, is expressed by several tissues and is known to activate B cells⁹⁶. CLCF-1 is secreted and forms heterodimers with either cytokine receptor-like factor 1 (CRLF1) or soluble ciliary neurotrophic receptor alpha (sCNTFR α), resulting in a composite cytokine⁹⁷. CLCF-1 was isolated by galactose affinity chromatography and found in the active fraction of plasma from patients with recurrent FSGS, at levels up to 100 times higher than the controls³⁶. CLCF-1 decreased nephrin expression in glomeruli and podocytes⁹⁶. The incubation of murine podocytes with CLCF-1 disrupted the actin cytoskeleton in time- and concentration-dependent manners and led to a motile podocyte phenotype⁹⁶. Similar to unfractionated FSGS serum, CLCF-1 isolated by galactose affinity chromatography induced albuminuria in mice and increased albumin permeability in isolated rat glomeruli through the activation of the JAK/STAT pathway, however, these experiments did not exclude the involvement of other galactose-binding permeability factors⁹⁶. This effect can be reversed by incubation with a monoclonal antibody against CRLF1 or with JAK/STAT inhibitors, suggesting that antibodies targeting CLCF-1 or its receptors and JAK2/STAT3 inhibitors may be novel therapeutic agents to treat FSGS⁹⁷. Recombinant human CRLF1 increased the albumin permeability in isolated rat glomeruli, which was inhibited by a heterodimer composed of CRLF1 and co-secreted molecule cytokine receptor-like factor 1⁹⁷.

Angiopietin like-4 (angptl-4)

Angptl-4 is a 45 to 65 kDa glycoprotein and acute phase reactant expressed in adipose tissue, liver, skeletal muscle, and the heart, and exists in two isoforms, with a high (pI>8) and a neutral (pI=7) isoelectric point, depending on the sialic acid content^{98,99}. A hyposialylated form secreted from podocytes can induce proteinuria in animal models, through an effect on either the GBM or endothelial cells,

and by reducing the anionic charges of the glomerular filtration barrier. In contrast, a neutral pI sialylated form is released into the circulation from other tissues and can reduce proteinuria by binding $\alpha\text{v}\beta 5$ integrins on the glomerular endothelium; it also induces hypertriglyceridemia via the inhibition of lipoprotein lipase. Rats overexpressing angptl-4 outside the kidney have high circulating levels of this factor but do not develop proteinuria, however, mice with podocyte overexpression of angptl-4 have normal circulating levels of the protein and do develop proteinuria¹⁰⁰. Glomerular angptl-4 expression and urinary angptl-4 levels also increased in diabetic rats¹⁰¹. In healthy individuals, angptl-4 is minimally expressed in the glomeruli, but is upregulated in the serum and urine of MCD patients¹⁰². Initial studies found that high circulating angptl-4 levels and glomerular expression coincided in human MCD and was related to proteinuria. However, subsequent studies have reported conflicting results.

The serum levels of angptl-4 were also elevated in patients with FSGS and MN compared with the controls, in addition to active MCD⁹⁸. In a recent study, Cara-Fuentes et al.¹⁰³ found that there was no correlation between serum angptl-4 levels and proteinuria in MCD, FSGS, and MN patients during relapse. Furthermore, the serum angptl-4 levels were decreased in MCD patients during relapse compared with the same patients in remission and the controls. In contrast with a previous study, no or minimal angptl-4 staining by immunofluorescence was observed in kidney tissue from MCD patients in relapse. They also found that urinary angptl-4 excretion in relapsed patients correlated with proteinuria in the FSGS and MN groups, however, this positive correlation was not found in MCD patients during relapse. Increased urinary angptl-4 excretion may be the result rather than the cause of proteinuria. In addition, no increased angptl-4 expression was observed in human podocytes cultured with sera from MCD patients in relapse compared with those exposed to sera from MCD patients in remission. Therefore, the clinical significance of angptl-4 in MCD remains to be determined given the contrasting findings between the relapse and remission groups. Nonetheless, angptl-4 remains a strong candidate for explaining MCD both in human and in experimental models. How sialylated angptl-4 could be used to treat NS is yet to be determined, however, off-target effects, such as

the inhibition of lipoprotein lipase leading to hypertriglyceridemia in NS, could be exacerbated^{98,100}. Previous studies have demonstrated that the modification of soluble angptl-4 sialylation or of key amino acids can be successfully used to treat proteinuria^{104,105}.

Interleukin-13 (IL-13)

IL-13 is a type of T cell-derived cytokine¹⁰⁶. IL-13 has been reported to be increased during MCD relapse, and the expression of the IL-13 gene is elevated both in CD4 and CD8 lymphocytes in children with steroid-sensitive NS during recurrences^{107,108}. This increase is associated with higher levels of IL-13 in the cytoplasm of T-cells¹⁰⁸ and the downregulation of IL-8 and IL-12 pro-inflammatory cytokines in monocytes¹⁰⁹. In addition, Wistar rats overexpressing IL-13 developed clinical NS and showed glomerular lesions identical to those observed in MCN, including an 80% effacement of podocyte foot processes via electron microscopy, the downregulation of nephrin, podocin and dystroglycan, and an increase in podocyte expression of CD80⁵⁷. We recently found pathological molecular changes in podocyte proteins by IL-13. We previously reported that IL-13 increased podocyte permeability through the modulation of ZO-1¹¹⁰, modulated the changes of cytoskeletal and adaptor proteins¹¹¹, and increased podocyte apoptosis¹¹², resulting in proteinuria and podocyte injury.

However, the supposed role of circulating IL-13 in proteinuria in MCD patients is questionable because the elevated serum IL-13 levels were not found in MCD patients during relapse⁶⁶ and have been reported in patients with NS during remission compared with the controls¹¹³. In addition, although serum IL-13 is increased in patients with asthma and with atopic dermatitis, only a few present associated proteinuria^{114,115}. Further experimental and clinical research on the patho-mechanism of IL-13 on the development of proteinuria and podocyte injury is needed.

Interleukin-8 (IL-8)

A pathogenic role for IL-8, a chemokine, in glomerular permeability was suggested after serum levels of IL-8 were

found to be increased in MCD in relapse compared with those observed during remission and in the controls^{116,117}. The urinary IL-8 level was 2.9-fold higher in NS patients in relapse compared to patients in remission, and levels had a positive correlation with individual proteinuria values. Urinary IL-8 was also significantly higher in relapsed steroid-sensitive NS patients than in steroid-sensitive NS patients in remission¹¹⁸. Rats infused with IL-8 that reached a serum IL-8 concentration similar to that observed in MCD patients developed mild proteinuria, and the administration of anti-IL-8-neutralizing antibody prevented proteinuria, suggesting a causal relationship between IL-8 and proteinuria¹¹⁹. In this model, IL-8 increased GBM glycosaminoglycan catabolism and reduced anionic charges at the GBM¹²⁰. However, rat podocytes incubated with high concentrations of human IL-8 did not show any alteration in permeability, heparan sulfate (HS) proteoglycan gene expression, or HS synthesis¹¹⁷. Unfortunately, further studies on the role of IL-8 for the development of proteinuria in NS have not been reported.

CD40

CD40 belongs to the TNF receptor superfamily and is expressed preferentially in B cells, monocytes/macrophages, dendritic cells, platelets, as well as non-immune cells, such as endothelial and smooth muscle cells, and is involved in the adaptive immune response¹²¹. The binding of CD40 to CD40 ligand expressed in T cells and platelets, that is, the CD40/CD40L complex, mediates pro-inflammatory events¹²². The CD40 ligand activates endothelium and leads to increased expression of chemokines, metalloproteases, uPA, and suPAR¹²³. CD40 is also constitutively expressed in podocytes, where activation by CD40L promotes the redistribution of nephrin and increases permeability to albumin in isolated glomeruli¹²³⁻⁵. CD40 was detected in the glomeruli and serum anti-CD40 antibodies were detected in recurrent FSGS patients, with levels which correlated with the post-transplant recurrence of the disease. As such, anti-CD40 antibodies may help to identify patients at high risk of post-transplant outcome¹²³. Even though anti-CD40 antibodies from recurrent FSGS sera did not detect recombinant human CD40, purified anti-CD40 antibodies from

recurrent FSGS sera were found to disrupt the podocyte actin cytoskeleton *in vitro*. This finding points to a post-translational modification of the CD40 molecule *in vivo*, which is necessary for detection with anti-CD40 antibodies¹²³. The authors also suggested that anti-CD40 antibodies play a pathogenetic role in the development of recurrent FSGS, potentially via its interaction with suPAR, facilitating or prolonging suPAR-mediated integrin activation, thus potentiating enhanced podocyte motility and disruption of the slit diaphragm¹²³. Injection with the CD40/anti-CD40 complex was found to induce proteinuria in mice, while infusion with anti-CD40L antibodies reduced proteinuria and the progression of adriamycin nephrosis^{126,127}. The size of IgG anti-CD40 antibodies is approximately 150 kDa⁹⁵, which contradicts previous findings where the active fraction of FSGS sera was found to be 30-50 kDa^{28,40}. Anti-CD40 blocking antibodies (ASKP1240 or lucatumumab) are already commercially available and could become potential treatment options to be tested in clinical trials⁴¹.

Heparanase

Heparanase, an endo- β (1-4)-D-glucuronidase that cleaves the glycosidic bond within the HS chain via hydrolysis, is involved in various biologic processes. Glomerular heparanase degrades HS polysaccharide side chains in GBM and thereby induces proteinuria via the loss of anionic charges, alterations in the glomerular cell-GBM interactions due to loss of HS, and the release of HS-bound factors, such as growth factors, cytokines, and chemokines, and bioactive HS fragments in glomeruli¹²⁸.

Heparanase has been found to play an important role of in several proteinuric diseases. A disturbance in heparanase expression was detected in the urine and glomeruli of proteinuric patients with MCD, FSGS, MN, and diabetic nephropathy¹²⁸⁻¹³². In steroid-sensitive NS in children, Holt et al.¹²⁹ found increased heparanase expression in the kidney glomeruli, increased heparanase concentration in the urine, and decreased heparanase concentration in the serum. A discrepancy between serum and urinary heparanase was associated with the accumulation of heparanase deposits along the vascular walls, the presence of inhibitors in the serum, or a non-proportional increase in heparanase versus

changes in proteinuria. The importance of heparanase has also been demonstrated in the several forms of experimental glomerular diseases. Increased heparanase expression and/or decreased HS was found in the early stages of experimental anti-GBM antibody disease¹³³, passive Heymann nephritis as an experimental MN model¹³⁴⁻¹³⁶, and puromycin aminonucleoside or adriamycin nephropathy^{137,138}. The expression of heparanase was found to remain stable over the course of these disorders.

However, the role of heparanase in the development of proteinuria has recently been challenged. A strong reduction in glycosaminoglycan-associated anionic sites in GBM without renal changes casts doubt on the primary role of HS in charge-selective filtration, suggesting that an increase in heparanase expression and a decrease in HS in GBM in glomerular diseases is a consequence rather than a cause of proteinuria^{128,139}. Indeed, proteinuria-lowering interventions, such as the inhibition of the renin-angiotensin-aldosterone system or reactive oxygen species scavengers, which are known to be unrelated to heparanase, also induced a regression of heparanase expression¹³⁸. Adriamycin-injected wild-type mice developed severe albuminuria, while

ADR-injected mice with transgenic heparanase overexpression exhibited only mild proteinuria and preserved podocyte phenotypes. Therefore, heparanase may have a nephroprotective role in adriamycin-induced NS, most likely via a mechanism independent of HS degradation¹⁴⁰. No further studies on the role of heparanase in the development of proteinuria have been carried out.

Conclusions

The biologic advances in our understanding of the pathogenesis of podocytopathies have facilitated the identification of various podocyte molecules and several candidate mediators that could play a role in FSGS or MCD. The pathogenic sequences of the permeability factors involved in the development of proteinuria and NS are presented in Fig. 1. Despite variations in the severity of correlation and the type of NS, permeability factors in MCD are VPF, hemopexin, CD80, angptl-4, heparanase, IL-13, and IL-8. On the other hand, suPAR, CLCF-1, and CD40 autoantibodies are permeability factors that would be more applicable in

Fig. 1. The pathophysiology of proteinuria and nephrotic syndrome.

FSGS. However, none of these molecules have been identified as the unique primary cause of either MCD or FSGS^{6,9,26,40,95}. Currently, new drugs are being developed that target podocyte molecules and pathogenic mediators involved in the pathogenesis of MCD/FSGS and recurrent FSGS.

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

Acknowledgments

This report was supported by the Clinical Research Fund of Chungbuk National University Hospital and Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (NRF-2013R1A1A4A03006207 and 2016R1D1A1B03931888).

ORCID

Tae-Sun Ha <http://orcid.org/0000-0002-1249-9342>

References

- Eddy AA, Symons JM. Nephrotic syndrome in childhood. *Lancet* 2003;362:629-39.
- Sinha A, Bagga A. Nephrotic syndrome. *Indian J Pediatr* 2012;79:1045-55.
- Vivarelli M, Massella L, Ruggiero B, Emma F. Minimal change disease. *Clin J Am Soc Nephrol* 2017;12:332-45.
- Noone DG, Iijima K, Parekh R. Idiopathic nephrotic syndrome in children. *Lancet* 2018;392(10141):61-74.
- Ha TS. Roles of adaptor proteins in podocyte biology. *World J Nephrol* 2013;2:1-10.
- Cara-Fuentes G, Clapp WL, Johnson RJ, Garin EH. Pathogenesis of proteinuria in idiopathic minimal change disease: molecular mechanisms. *Pediatr Nephrol* 2016;31:2179-89.
- Ha TS. Genetics of hereditary nephrotic syndrome: a clinical review. *Korean J Pediatr* 2017;60:55-63.
- Shalhoub RJ. Pathogenesis of lipoid nephrosis: a disorder of T-cell function. *Lancet* 1974;2:556-60.
- Davin JC. The glomerular permeability factors in idiopathic nephrotic syndrome. *Pediatr Nephrol* 2016;31:207-15.
- Musante L, Candiano G, Zennaro C, Bruschi M, Carraro M, Artero M, et al. Humoral permeability factors in the nephrotic syndrome: a compendium and prospectus. *J Nephrol* 2001;14 Suppl 4:S48-50.
- Laguerre G, Branellec A, Niaudet P, Heslan JM, Guillot F, Lang P. Transmission of nephrotic syndrome to two neonates. Spontaneous regression. *Presse Med* 1991;20:255-7.
- Kemper MJ, Wolf G, Müller-Wiefel DE. Transmission of glomerular permeability factor from a mother to her child. *N Engl J Med* 2001;344:386-7.
- Zimmerman SW. Increased urinary protein excretion in the rat produced by serum from a patient with recurrent focal glomerular sclerosis after renal transplantation. *Clin Nephrol* 1984;22: 32-8.
- Sharma M, Sharma R, Reddy SR, McCarthy ET, Savin VJ. Proteinuria after injection of human focal segmental glomerulosclerosis factor. *Transplantation* 2002;73: 366-72.
- Raj L, Hoyer JR, Michael AF. Steroid-resistant nephrotic syndrome: recurrence after transplantation. *Ann Intern Med* 1972;77:581-6.
- Hoyer JR, Vernier RL, Najarian JS, Raj L, Simmons RL, Michael AF. Recurrence of idiopathic nephrotic syndrome after renal transplantation. *Lancet* 1972;2:343-8.
- Fine RN. Recurrence of nephrotic syndrome/focal segmental glomerulosclerosis following renal transplantation in children. *Pediatr Nephrol* 2007;22:496-502.
- Zimmerman SW. Plasmapheresis and dipyridamole for recurrent focal glomerular sclerosis. *Nephron* 1985;40:241-5.
- Artero M, Biava C, Amend W, Tomlanovich S, Vincenti F. Recurrent focal glomerulosclerosis: Natural history and response to therapy. *Am J Med* 1992;92:375-83.
- Dantal J, Bigot E, Bogers W, Testa A, Kriaa F, Jacques Y, et al. Effect of plasma protein adsorption on protein excretion in kidney-transplant recipients with recurrent nephrotic syndrome. *N Engl J Med* 1994;330:7-14.
- Artero ML, Sharma R, Savin VJ, Vincenti F. Plasmapheresis reduces proteinuria and serum capacity to injure glomeruli in patients with recurrent focal glomerulosclerosis. *Am J Kidney Dis* 1994; 23:574-81.
- Gohh RY, Yango AF, Morrissey PE, Monaco AP, Gautam A, Sharma M, et al. Preemptive plasmapheresis and recurrence of FSGS in high-risk renal transplant recipients. *Am J Transplant* 2005;5:2907-12.
- Suthar K, Vanikar AV, Trivedi HL. Renal transplantation in primary focal segmental glomerulosclerosis using a tolerance induction protocol. *Transplant Proc* 2008;40:1108-10.
- Gallon L, Leventhal J, Skaro A, Kanwar Y, Alvarado A. Resolution of recurrent focal segmental glomerulosclerosis after retransplantation. *N Engl J Med* 2012;366:1648-9.
- Rea R, Smith C, Sandhu K, Kwan J, Tomson C. Successful transplant

- of a kidney with focal segmental glomerulosclerosis. *Nephrol Dial Transplant* 2001;16:416-7.
26. Maas RJ, Deegens JK, Wetzels JF. Permeability factors in idiopathic nephrotic syndrome: historical perspectives and lessons for the future. *Nephrol Dial Transplant* 2014;29:2207-16.
 27. Lagrue G, Xheneumont S, Branellec A, Hirbec G, Weil B. A vascular permeability factor elaborated from lymphocytes. I. Demonstration in patients with nephrotic syndrome. *Biomedicine* 1975; 23:37-40.
 28. Savin VJ, Sharma R, Sharma M, et al. Circulating factor associated with increased glomerular permeability to albumin in recurrent focal segmental glomerulosclerosis. *N Engl J Med* 1996;334: 878-83.
 29. Kachurina N, Chung CF, Benderoff E, Babayeva S, Bitzan M, Goodyer P, et al. Novel unbiased assay for circulating podocyte-toxic factors associated with recurrent focal segmental glomerulosclerosis. *Am J Physiol Renal Physiol* 2016;310:F1148-56.
 30. Lennon R, Singh A, Welsh GI, Coward RJ, Satchell S, Ni L, et al. Hemopexin induces nephrin-dependent reorganization of the actin cytoskeleton in podocytes. *J Am Soc Nephrol* 2008;19:2140-9.
 31. Bakker WW, Donker RB, Timmer A, van Pampus MG, van Son WJ, Aarnoudse JG, et al. Plasma hemopexin activity in pregnancy and preeclampsia. *Hypertens Pregnancy* 2007;26:227-39.
 32. Savin VJ, Sharma R, Lovell HB, Welling DJ. Measurement of albumin reflection coefficient with isolated rat glomeruli. *J Am Soc Nephrol* 1992;3:1260-9.
 33. Desideri S, Onions KL, Qiu Y, Ramnath RD, Butler MJ, Neal CR, et al. A novel assay provides sensitive measurement of physiologically relevant changes in albumin permeability in isolated human and rodent glomeruli. *Kidney Int* 2018;93:1086-97.
 34. Savin VJ, McCarthy ET, Sharma R, Charba D, Sharma M. Galactose binds to focal segmental glomerulosclerosis permeability factor and inhibits its activity. *Transl Res* 2008;151:288-92.
 35. Wang L, Tao T, Su W, Yu H, Yu Y, Qin J. A disease model of diabetic nephropathy in a glomerulus-on-a-chip microdevice. *Lab Chip* 2017;17:1749-60.
 36. Koyama A, Fujisaki M, Kobayashi M, Igarashi M, Narita M. A glomerular permeability factor produced by human T cell hybridomas. *Kidney Int* 1991;40:453-60.
 37. Cheung PK, Klok PA, Baller JF, Bakker WW. Induction of experimental proteinuria in vivo following infusion of human plasma hemopexin. *Kidney Int* 2000;57:1512-20.
 38. Sharma M, Zhou J, Gauchat JF, Sharma R, McCarthy ET, Srivastava T, et al. Janus kinase 2/signal transducer and activator of transcription 3 inhibitors attenuate the effect of cardiotrophin-like cytokine factor 1 and human focal segmental glomerulosclerosis serum on glomerular filtration barrier. *Transl Res* 2015;166:384-98.
 39. McCarthy ET, Sharma M, Savin VJ. Circulating permeability factors in idiopathic nephrotic syndrome and focal segmental glomerulosclerosis. *Clin J Am Soc Nephrol* 2010;5:2115-21.
 40. Savin VJ, McCarthy ET, Sharma M. Permeability factors in nephrotic syndrome and focal segmental glomerulosclerosis. *Kidney Res Clin Pract* 2012;31:205-13.
 41. Wada T, Nangaku M. A circulating permeability factor in focal segmental glomerulosclerosis: the hunt continues. *Clin Kidney J* 2015;8:708-15.
 42. Bakker WW, van der LSM, Vos JT, Hoedemaeker PJ. The glomerular polyanion (GPA) of the rat kidney. I. Concanavalin-A activated cells affect the glomerular polyanion in vitro. *Nephron* 1982;31: 68-74.
 43. Bakker WW, van LWH, Hene RJ, Desmit EM, van der HGK, Vos JT. Loss of glomerular polyanion in vitro induced by mononuclear blood cells from patients with minimal-change nephrotic syndrome. *Am J Nephrol* 1986;6:107-11.
 44. Boulton-Jones JM, Tulloch I, Dore B, McLay A. Changes in the glomerular capillary wall induced by lymphocyte products and serum of nephrotic patients. *Clin Nephrol* 1983;20:72-7.
 45. Yoshizawa N, Kusumi Y, Matsumoto K, Oshima S, Takeuchi A, Kawamura O, et al. Studies of a glomerular permeability factor in patients with minimal-change nephrotic syndrome. *Nephron* 1989;51:370-6.
 46. Bakker WW, van Dael CM, Pierik LJ, van Wijk JA, Nauta J, Borghuis T, et al. Altered activity of plasma hemopexin in patients with minimal change disease in relapse. *Pediatr Nephrol* 2005;20: 1410-5.
 47. Cheung PK, Klok PA, Bakker WW. Minimal change-like glomerular alterations induced by a human plasma factor. *Nephron* 1996;74: 586-93.
 48. Shimada M, Ishimoto T, Lee PY, Lanaspa MA, Rivard CJ, Roncal-Jimenez CA, et al. Toll-like receptor 3 ligands induce CD80 expression in human podocytes via an NF- κ B-dependent pathway. *Nephrol Dial Transplant* 2012;27:81-9.
 49. Wing K, Onishi Y, Prieto-Martin P, Yamaguchi T, Miyara M, Fehervari Z, et al. CTLA-4 control over Foxp3+ regulatory T cell function. *Science* 2008;322:271-5.
 50. Mosser DM, Zhang X. Interleukin-10: new perspectives on an old cytokine. *Immunol Rev* 2008;226:205-18.
 51. Shimada M, Araya C, Rivard C, Ishimoto T, Johnson RJ, Garin EH. Minimal change disease: a "two-hit" podocyte immune disorder? *Pediatr Nephrol* 2011;26:645-9.
 52. Cara-Fuentes G, Wasserfall CH, Wang H, Johnson RJ, Garin EH. Minimal change disease: a dysregulation of the podocyte CD80-CTLA-4 axis? *Pediatr Nephrol* 2014;29:2333-40.
 53. Ohl K, Eberhardt C, Spink C, Zahn K, Wagner N, Eggermann T, et al. CTLA4 polymorphisms in minimal change nephrotic syndrome in children: a case-control study. *Am J Kidney Dis* 2014;63: 1074-5.
 54. Spink C, Stege G, Tenbrock K, Harendza S. The CTLA-4+49GG genotype is associated with susceptibility for nephrotic kidney diseases. *Nephrol Dial Transplant* 2013;28:2800-5.
 55. Cara-Fuentes G, Wei C, Segarra A, Ishimoto T, Rivard C, Johnson RJ, et al. CD80 and suPAR in patients with minimal change disease and focal segmental glomerulosclerosis: diagnostic and

- pathogenic significance. *Pediatr Nephrol* 2014;29:1363-71.
56. Reiser J, Mundel P. Danger signaling by glomerular podocytes defines a novel function of inducible B7-1 in the pathogenesis of nephrotic syndrome. *J Am Soc Nephrol* 2004;15:2246-8.
 57. Lai KW, Wei CL, Tan LK, Tan PH, Chiang GS, Lee CG, et al. Overexpression of interleukin-13 induces minimal-change-like nephropathy in rats. *J Am Soc Nephrol* 2007;18:1476-85.
 58. Reiser J, von Gersdorff G, Loos M, Oh J, Asanuma K, Giardino L, et al. Induction of B7-1 in podocytes is associated with nephrotic syndrome. *J Clin Invest* 2004;113:1390-7.
 59. Khullar B, Balyan R, Oswal N, Jain N, Sharma A, Abdin MZ, et al. Interaction of CD80 with Neph1: a potential mechanism of podocyte injury. *Clin Exp Nephrol* 2018;22:508-16.
 60. Ishimoto T, Shimada M, Gabriela G, Kosugi T, Sato W, Lee PY, et al. Toll-like receptor 3 ligand, polyIC, induces transient proteinuria and glomerular CD80, and increases urinary CD80 in mice. *Nephrol Dial Transplant* 2013;28:1439-46.
 61. Garin EH, Diaz LN, Mu W, Wasserfall C, Araya C, Segal M, et al. Urinary CD80 excretion increases in idiopathic minimal-change disease. *J Am Soc Nephrol* 2009;20:260-6.
 62. Garin EH, Mu W, Arthur JM, Rivard CJ, Araya CE, Shimada M, et al. Urinary CD80 is elevated in minimal change disease but not in focal segmental glomerulosclerosis. *Kidney Int* 2010;78:296-302.
 63. Mishra OP, Kumar R, Narayan G, Srivastava P, Abhinay A, Prasad R. Toll-like receptor 3 (TLR-3), TLR-4 and CD80 expression in peripheral blood mononuclear cells and urinary CD80 levels in children with idiopathic nephrotic syndrome. *Pediatr Nephrol* 2017;32:1355-61.
 64. Fiorina P, Vergani A, Bassi R, Niewczas MA, Altintas MM, Pezzolesi MG. Role of podocyte B7-1 in diabetic nephropathy. *J Am Soc Nephrol* 2014;25:1415-29.
 65. Yu CC, Fornoni A, Weins A, Hakrrouch S, Maiquel D, Sageshima J, et al. Abatacept in B7-1-positive proteinuric kidney disease. *N Engl J Med* 2013;369:2416-23.
 66. Ishimoto T, Cara-Fuentes G, Wang H, Shimada M, Wasserfall CH, Winter WE, et al. Serum from minimal change patients in relapse increases CD80 expression in cultured podocytes. *Pediatr Nephrol* 2013;28:1803-12.
 67. Garin EH, Reiser J, Cara-Fuentes G, Wei C, Matar D, Wang H, et al. Case series: CTLA4-IgG1 therapy in minimal change disease and focal segmental glomerulosclerosis. *Pediatr Nephrol* 2015;30:469-77.
 68. Delville M, Baye E, Durrbach A, Audard V, Kofman T, Braun L. B7-1 blockade does not improve post-transplant nephrotic syndrome caused by recurrent FSGS. *J Am Soc Nephrol* 2016;27:2520-7.
 69. Estreicher A, Mühlhauser J, Carpentier JL, Orci L, Vassalli JD. The receptor for urokinase type plasminogen activator polarizes expression of the protease to the leading edge of migrating monocytes and promotes degradation of enzyme inhibitor complexes. *J Cell Biol* 1990;111:783-92.
 70. Nykjaer A, Møller B, Todd RF 3rd, Christensen T, Andreasen PA, Gliemann J, et al. Urokinase receptor. An activation antigen in human T lymphocytes. *J Immunol* 1994;152:505-16.
 71. Maas RJ, Deegens JK, Wetzels JF. Serum suPAR in patients with FSGS: trash or treasure? *Pediatr Nephrol* 2013;28:1041-8.
 72. Roldan AL, Cubellis MV, Masucci MT, Behrendt N, Lund LR, Dano K, et al. Cloning and expression of the receptor for human urokinase plasminogen activator, a central molecule in cell surface, plasmin dependent proteolysis. *EMBO J* 1990;9:467-74.
 73. Smith HW, Marshall CJ. Regulation of cell signalling by uPAR. *Nat Rev Mol Cell Biol* 2010;11:23-36.
 74. Dass K, Ahmad A, Azmi AS, Sarkar SH, Sarkar FH. Evolving role of uPA/uPAR system in human cancers. *Cancer Treat Rev* 2008;34:122-36.
 75. Wei C, El Hindi S, Li J, Fornoni A, Goes N, Sageshima J, et al. Circulating urokinase receptor as a cause of focal segmental glomerulosclerosis. *Nat Med* 2011;17:952-60.
 76. Kreidberg JA, Donovan MJ, Goldstein SL, Rennke H, Shepherd K, Jones RC, et al. Alpha 3 beta 1 integrin has a crucial role in kidney and lung organogenesis. *Development* 1996;122:3537-47.
 77. Wei C, Moller CC, Altintas MM, Li J, Schwarz K, Zaccogna S, et al. Modification of kidney barrier function by the urokinase receptor. *Nat Med* 2008;14:55-63.
 78. Wei C, Trachtman H, Li J, Dong C, Friedman AL, Gassman JJ, et al. Circulating suPAR in two cohorts of primary FSGS. *J Am Soc Nephrol* 2012;23:2051-9.
 79. Li F, Zheng C, Zhong Y, Zeng C, Xu F, Yin R, et al. Relationship between serum soluble urokinase plasminogen activator receptor level and steroid responsiveness in FSGS. *Clin J Am Soc Nephrol* 2014;9:1903-11.
 80. Huang J, Liu G, Zhang YM, Cui Z, Wang F, Liu XJ, et al. Plasma soluble urokinase receptor levels are increased but do not distinguish primary from secondary focal segmental glomerulosclerosis. *Kidney Int* 2013;84:366-72.
 81. Meijers B, Maas RJ, Sprangers B, Claes K, Poesen R, Bammens B, et al. The soluble urokinase receptor is not a clinical marker for focal segmental glomerulosclerosis. *Kidney Int* 2014;85:636-40.
 82. Wada T, Nangaku M, Maruyama S, Imai E, Shoji K, Kato S, et al. A multicenter cross-sectional study of circulating soluble urokinase receptor in Japanese patients with glomerular disease. *Kidney Int* 2014;85:641-8.
 83. Pinale JM, Mariani LH, Kapoor S, Zhang J, Weyant R, Song PX, et al. Nephrotic Syndrome Study Network () A reassessment of soluble urokinase-type plasminogen activator receptor in glomerular disease. *Kidney Int* 2015;87:564-74.
 84. Maas RJ, Wetzels JF, Deegens JK. Serum-soluble urokinase receptor concentration in primary FSGS. *Kidney Int* 2012;81:1043-4.
 85. Bock ME, Price HE, Gallon L, Langman CB. Serum soluble urokinase-type plasminogen activator receptor levels and idiopathic FSGS in children: a single-center report. *Clin J Am Soc Nephrol* 2013;8:1304-11.
 86. Sinha A, Bajpai J, Saini S, Bhatia D, Gupta A, Puraswani M, et al. Serum-soluble urokinase receptor levels do not distinguish focal

- segmental glomerulosclerosis from other causes of nephrotic syndrome in children. *Kidney Int* 2014;85:649-58.
87. Harita Y, Ishizuka K, Tanego A, Sugawara N, Chikamoto H, Akioka Y, et al. Decreased glomerular filtration as the primary factor of elevated circulating suPAR levels in focal segmental glomerulosclerosis. *Pediatr Nephrol* 2014;29:1553-60.
 88. Huang J, Liu G, Zhang YM, Cui Z, Wang F, Liu XJ, et al. Urinary soluble urokinase receptor levels are elevated and pathogenic in patients with primary focal segmental glomerulosclerosis. *BMC Med* 2014;12:81.
 89. Wei C, Li J, Adair BD, Zhu K, Cai J, Merchant M, et al. uPAR isoform 2 forms a dimer and induces severe kidney disease in mice. *J Clin Invest* 2019 [Epub ahead of print]
 90. Alfano M, Cinque P, Giusti G, Proietti S, Nebuloni M, Danese S, et al. Full-length soluble urokinase plasminogen activator receptor down-modulates nephrin expression in podocytes. *Sci Rep* 2015;5:13647.
 91. Delville M, Sigdel TK, Wei C, Li J, Hsieh SC, Fornoni A, et al. A circulating antibody panel for pretransplant prediction of FSGS recurrence after kidney transplantation. *Sci Transl Med* 2014;6:256ra136.
 92. Cathelin D, Placier S, Ploug M, Verpont MC, Vandermeersch S, Luque Y, et al. Administration of recombinant soluble urokinase receptor per se is not sufficient to induce podocyte alterations and proteinuria in mice. *J Am Soc Nephrol* 2014;25:1662-8.
 93. Segarra-Medrano A, Carnicer-Cáceres C, Arbós-Via MA, Quiles-Pérez MT, Agraz-Pamplona I, Ostos-Roldán E. Biological markers of nephrotic syndrome: a few steps forward in the long way. *Nefrologia* 2012;32:558-72.
 94. Kronbichler A, Saleem MA, Meijers B, Shin JI. Soluble urokinase receptors in focal segmental glomerulosclerosis: a review on the scientific point of view. *J Immunol Res* 2016;2068691.
 95. Königshausen E, Sellin L. Circulating permeability factors in primary focal segmental glomerulosclerosis: A review of proposed candidates. *Biomed Res Int* 2016;2016:3765608.
 96. Savin VJ, Sharma M, Zhou J, Gennochi D, Fields T, Sharma R, et al. Renal and hematological effects of CLCF-1, a B-cell-stimulating cytokine of the IL-6 family. *J Immunol Res* 2015;2015:714964.
 97. Sharma M, Zhou J, Gauchat JF, Sharma R, McCarthy ET, Srivastava T, et al. Janus kinase 2/signal transducer and activator of transcription 3 inhibitors attenuate the effect of cardiotrophin-like cytokine factor 1 and human focal segmental glomerulosclerosis serum on glomerular filtration barrier. *Transl Res* 2015;166:384-98.
 98. Clement LC, Macé C, Avila-Casado C, Joles JA, Kersten S, Chugh SS. Circulating angiopoietin-like 4 links proteinuria with hypertriglyceridemia in nephrotic syndrome. *Nat Med* 2014;20:37-46.
 99. Bertelli R, Bonanni A, Caridi G, Canepa A, Ghiggeri GM. Molecular and cellular mechanisms for proteinuria in minimal change disease. *Front Med (Lausanne)* 2018;5:170.
 100. Chugh SS, Clement LC, Mace C. New insights into human minimal change disease: lessons from animal models. *Am J Kidney Dis* 2012;59:284-92.
 101. Ma J, Chen X, Li JS, Peng L, Wei SY, Zhao SL, et al. Upregulation of podocyte-secreted angiopoietin-like-4 in diabetic nephropathy. *Endocrine* 2015;49:373-84.
 102. Clement LC, Avila-Casado C, Macé C, Soria E, Bakker WW, Kersten S, et al. Podocyte-secreted angiopoietin-like-4 mediates proteinuria in glucocorticoid-sensitive nephrotic syndrome. *Nat Med* 2011;17:117-22.
 103. Cara-Fuentes G, Segarra A, Silva-Sanchez C, Wang H, Lanaspá MA, Johnson RJ, et al. Angiopoietin-like-4 and minimal change disease. *PLoS One* 2017;12:e0176198.
 104. Macé C, Chugh SS. Nephrotic syndrome: components, connections, and angiopoietin-like 4-related therapeutics. *J Am Soc Nephrol* 2014;25:2393-8.
 105. Chugh SS, Macé C, Clement LC, Del Noga Avila M, Marshall CB. Angiopoietin-like 4 based therapeutics for proteinuria and kidney disease. *Front Pharmacol* 2014;5:23.
 106. McKenzie AN, Culpepper JA, de Waal Malefyt R, Briere F, Punnonen J, Aversa G, et al. Interleukin 13, a T-cell-derived cytokine that regulates human monocyte and B-cell function. *Proc Natl Acad Sci USA* 1993;90:3735-40.
 107. Yap HK, Cheung W, Murugasu B, Sim SK, Seah CC, Jordan SC. Th1 and Th2 cytokine mRNA profiles in childhood nephrotic syndrome: evidence for increased IL-13 mRNA expression in relapse. *J Am Soc Nephrol* 1999;10:529-37.
 108. Cheung W, Wei CL, Seah CC, Jordan SC, Yap HK. Atopy, serum IgE, and interleukin-13 in steroid-responsive nephrotic syndrome. *Pediatr Nephrol* 2004;19:627-32.
 109. Chen SP, Cheung W, Heng CK, Jordan SC, Yap HK. Childhood nephrotic syndrome in relapse is associated with down-regulation of monocyte CD14 expression and lipopolysaccharide-induced tumour necrosis factor- α production. *Clin Exp Immunol* 2003;134:111-9.
 110. Park SJ, Saleem MA, Nam JA, Ha TS, Shin JI. Effects of interleukin-13 and montelukast on the expression of zonula occludens-1 in human podocytes. *Yonsei Med J* 2015;56:426-32.
 111. Ha TS, Nam JA, Seong SB, Saleem MA, Park SJ, Shin JI. Montelukast improves the changes of cytoskeletal and adaptor proteins of human podocytes by interleukin-13. *Inflamm Res* 2017;66:793-802.
 112. Lee KH, Oh JY, Seong SB, Ha TS, Shin JI. Interleukin-13 increases podocyte apoptosis in cultured human podocytes. *Child Kidney Dis* 2018;22:22-7.
 113. Kanai T, Shiraishi H, Yamagata T, Ito T, Odaka J, Saito T, et al. Th2 cells predominate in idiopathic steroid-sensitive nephrotic syndrome. *Clin Exp Nephrol* 2010;14:578-83.
 114. Bagnasco D, Ferrando M, Varricchi G, Passalacqua G, Canonica GW. A critical evaluation of anti-IL-13 and anti-IL-4 strategies in severe asthma. *Int Arch Allergy Immunol* 2016;170:122-31.
 115. Matsunaga MC, Yamauchi PS. IL-4 and IL-13 inhibition in atopic dermatitis. *J Drugs Dermatol* 2016;15:925-9.
 116. Garin EH, Blanchard DK, Matsushima K, Djeu JY. IL-8 production by peripheral blood mononuclear cells in nephrotic patients.

- Kidney Int 1994;45:1311-7.
117. Cho MH, Lee HS, Choe BH, Kwon SH, Chung KY, Koo JH, et al. Interleukin-8 and tumor necrosis factor-alpha are increased in minimal change disease but do not alter albumin permeability. *Am J Nephrol* 2003;23:260-6.
118. Souto MF, Teixeira AL, Russo RC, Penido MG, Silveira KD, Teixeira MM, et al. Immune mediators in idiopathic nephrotic syndrome: evidence for a relation between interleukin 8 and proteinuria. *Pediatr Res* 2008;64:637-42.
119. Garin EH, Laflam P, Chandler L. Anti-interleukin 8 antibody abolishes effects of lipoid nephrosis cytokine. *Pediatr Nephrol* 1998;12:381-5.
120. Garin EH, West L, Zheng W. Effect of interleukin-8 on glomerular sulfated compounds and albuminuria. *Pediatr Nephrol* 1997;11:274-9.
121. Quezada SA, Jarvinen LZ, Lind EF, Noelle RJ. CD40/CD154 interactions at the interface of tolerance and immunity. *Annu Rev Immunol* 2004;22:307-28.
122. Hassan GS, Merhi Y, Mourad W. CD40 ligand: a neo-inflammatory molecule in vascular diseases. *Immunobiology* 2012;217:521-32.
123. Delville M, Sigdel TK, Wei C, Li J, Hsieh SC, Fornoni A, et al. A circulating antibody panel for pretransplant prediction of FSGS recurrence after kidney transplantation. *Sci Transl Med* 2014;6:256ra136.
124. Kairaitis L, Wang Y, Zheng L, Tay YC, wang YD, Harris C. Blockade of CD40-CD40 ligand protects against renal injury in chronic proteinuric renal disease. *Kidney Int* 2003;64:1265-72.
125. Doublier S, Zennaro C, Musante L, Spatola T, Candiano G, Bruschi M. Soluble CD40 ligand directly alters glomerular permeability and may act as a circulating permeability factor in FSGS. *PLoS One* 2017;12:e0188045.
126. Lee VW, Qin X, Wang Y, Zheng G, Wang Y, Wang Y. The CD40-CD154 co-stimulation pathway mediates innate immune injury in adriamycin nephrosis. *Nephrol Dial Transplant* 2010;25:717-30.
127. Kairaitis L, Wang Y, Zheng L, Tay YC, Wang Y, Harris DC. Blockade of CD40-CD40 ligand protects against renal injury in chronic proteinuric renal disease. *Kidney Int* 2003;64:1265-72.
128. van den Hoven MJ, Rops AL, Vlodavsky I, Levidiotis V, Berden JH, van der Vlag J. Heparanase in glomerular diseases. *Kidney Int* 2007;72:543-8.
129. Holt RC, Webb NJ, Ralph S, Davies J, Short CD, Brenchley PE. Heparanase activity is dysregulated in children with steroid-sensitive nephrotic syndrome. *Kidney Int* 2005;67:122-9.
130. Katz A, Van-Dijk DJ, Aingorn H, Erman A, Davies M, Darmon D, et al. Involvement of human heparanase in the pathogenesis of diabetic nephropathy. *Isr Med Assoc J* 2002;4:996-1002.
131. van den Hoven MJ, Rops AL, Bakker MA, Aten J, Rutjes N, Roestenberg P, et al. Increased expression of heparanase in overt diabetic nephropathy. *Kidney Int* 2006;70:2100-8.
132. Szymczak M, Kuźniar J, Klinger M. The role of heparanase in diseases of the glomeruli. *Arch Immunol Ther Exp (Warsz)* 2010;58:45-56.
133. Levidiotis V, Freeman C, Tikellis C, Cooper ME, Power DA. Heparanase inhibition reduces proteinuria in a model of accelerated anti-glomerular basement membrane antibody disease. *Nephrology (Carlton)* 2005;10:167-73.
134. Raats CJ, Luca ME, Bakker MA, Van Der Wal A, Heeringa P, Van Goor H, et al. Reduction in glomerular heparin sulfate correlates with complement deposition and albuminuria in active Heymann nephritis. *J Am Soc Nephrol* 1999;10:1689-99.
135. Levidiotis V, Freeman C, Tikellis C, Cooper ME, Power DA. Heparanase is involved in the pathogenesis of proteinuria as a result of glomerulonephritis. *J Am Soc Nephrol* 2004;15:68-78.
136. Levidiotis V, Freeman C, Punler M, Martinello P, Creese B, Ferro V, et al. A synthetic heparanase inhibitor reduces proteinuria in passive Heymann nephritis. *J Am Soc Nephrol* 2004;15:2882-92.
137. Levidiotis V, Kanellis J, Ierino FL, Power DA. Increased expression of heparanase in puromycin aminonucleoside nephrosis. *Kidney Int* 2001;60:1287-96.
138. Kramer A, van den Hoven M, Rops A, Wijnhoven T, van den Heuvel L, Lensen J, et al. Induction of glomerular heparanase expression in rats with adriamycin nephropathy is regulated by reactive oxygen species and the renin-angiotensin system. *J Am Soc Nephrol* 2006;17:2513-20.
139. van den Hoven MJ, Wijnhoven TJ, Li JP, Zcharia E, Dijkman HB, Wismans RG, et al. Reduction of anionic sites in the glomerular basement membrane by heparanase does not lead to proteinuria. *Kidney Int* 2008;73:278-87.
140. Assady S, Alter J, Axelman E, Zohar Y, Sabo E, Litvak M, et al. Nephroprotective effect of heparanase in experimental nephrotic syndrome. *PLoS One* 2015;10:e0119610.