Pediatric Acute Kidney Injury: Focusing on Diagnosis and Management

Myung Hyun Cho, M.D.

Department of Pediatrics, Hallym
University Sacred Heart Hospital,
Anyang, Republic of Korea

Corresponding author:

Myung Hyun Cho, M.D.
Address: Department of Pediatrics, Hallym
University Sacred Heart Hospital, 22
Gwanpyeong-ro 170 beon-gil, Dongan-
gu, Anyang 14068, Korea
Tel: +82-31-380-3730
Fax: +82-31-380-1900
E-mail: dong82dong82@naver.com

Received: 6 March 2020
Revised: 1 April 2020
Accepted: 9 April 2020

Acute kidney injury (AKI) is common in critically ill children, and is associated with increased mortality and long-term renal sequelae. The definition of pediatric AKI was standardized based on elevation in serum creatinine levels or decrease in urine output; accordingly, epidemiological studies have ensued. Although new biomarkers appear to detect AKI earlier and predict prognosis more accurately than traditional markers, they are not frequently used in clinical setting. There is no validated pharmacological intervention for AKI, so prevention and early detection are the mainstays of treatment. For high risk or early stage AKI patients, optimization of volume status and blood pressure, avoidance of nephrotoxins, and sufficient nutritional support are necessary, and have been demonstrated to be effective in preventing the occurrence of AKI and improving prognosis. Nevertheless, renal replacement therapy is needed when conservative care fails.

Key words: Acute kidney injury, Critical care, Pediatrics

Introduction

Acute kidney injury (AKI) is characterized by an abrupt deterioration of kidney function, and is common in critically ill children and adults. It occurs in approximately 30% of pediatric intensive care unit (PICU)¹⁾. Pediatric AKI has been associated with higher morbidity and mortality after adjustment for other risk factors¹⁾, and is a risk factor for hypertension and chronic kidney disease (CKD) in the long term^{2,3)}. Recently, many studies have been conducted in the field of pediatric AKI following adult studies and have prompted new interest. This review summarizes pediatric AKI, with a focus on diagnosis and management.

Definition of AKI

In 2012, the Kidney Disease: Improving Global Outcomes (KDIGO) criteria were established, and provided a standardized definition of AKI in children and adults⁴⁾. It was characterized by elevation in serum creatinine levels and/or decrease in urine output, based on the previously well-known, Risk, Injury, Failure; Loss, End-Stage Renal Disease (RIFLE) and Acute Kidney Injury Network (AKIN) criteria (Table 1)⁴⁾. The severity of AKI is staged according

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

to the amplitude of serum creatinine elevation from baseline value or the duration of compromised urine output.

Physicians sometimes, however, experience difficulty in diagnosing and staging AKI using the KDIGO definition. Many patients exhibit elevated serum creatinine levels without previous measurement. It is not integrated to set baseline value without available measurement data within 48 h, as specified by the KDIGO criteria. Some have used the lowest serum creatinine level measured before or after AKI as a baseline value. Others propose that the baseline serum creatinine level can calculate backward using an estimated glomerular filtration rate (eGFR) of 100–120 mL/min/1.73 m², assuming that previous kidney function was normal⁵⁾.

For neonates, the neonatal modified KDIGO definition was used in a recent large cohort (Table 1)^{6,7)}. Although it is based on a modification of the KDIGO definition, it is different with urinary output checked in 24 h blocks. Moreover, the baseline value is the lowest previous serum creatinine level because serum creatinine, which reflects maternal creatinine in the first few days, declines physiologically within weeks of life at a rate that varies with gestational age^{8,9)}.

AKI biomarkers

Serum creatinine is currently widely used as a biomarker for AKI; however, it has some limitations. It is insensitive

to small changes in GFR, and is not a real-time indicator. It may not change until up to 50% of kidney function is lost¹⁰⁾, and rise up to 72 h after an insult. Furthermore, its concentration is affected by age, sex, muscle mass, and volume status¹¹⁾. For two decades, many researchers have been searching new biomarkers that are rapid, sensitive, specific, inexpensive, noninvasive, and unaffected by clinical factors¹²⁾. Cystatin C, neutrophil gelatinase-associated lipocalin (NGAL), kidney injury molecule-1, N-acetyl-β-D-glucosaminidase, interleukin-18, liver-type fatty acid-binding protein, cycle arrest markers, and BPI containing family A member 2 have been evaluated. These appear to detect AKI earlier and predict prognosis more accurately than serum creatinine levels¹²⁾.

Serum cystatin C is produced by all nucleated cells and, is therefore, unaffected by clinical factors¹³⁾. Its level rises earlier (12–24 h after insult) than that of serum creatinine¹⁴⁾. In studies involving specific pediatric populations (children who underwent cardiac surgery, treated with aminoglycoside or contrast agent, and preterm infants with respiratory distress syndrome), it predicted AKI earlier than serum creatinine level, predicted persistent AKI, and showed greater sensitivity and specificity¹⁴⁻¹⁷⁾. Its disadvantage is that it is affected by high-dose corticosteroids, inflammation, and systemic diseases¹⁸⁾. NGAL, a proximal tubular protein, is the most rapid predictive biomarker, and is elevated <2 h after insult¹⁹⁾. Studies investigated NGAL in pediatrics have mainly involved patient undergoing cardiac surgery, and have reported that elevation of urinary NGAL

Table 1. Definitions and Staging of Kidney Disease: Improving Global Outcomes (KDIGO) and Neonatal Modified KDIGO Criteria for Acute Kidney Injury

Stage	Pediatric KDIGO criteria		Neonatal modified KDIGO criteria	
	Serum creatinine	Urine output	Serum creatinine	Urine output
1	1.5–1.9 times baseline within 7 days OR ≥0.3 mg/dL increase within 48 h	<0.5 mL/kg/h for 6–12 h	1.5–1.9 times baseline* within 7 days OR ≥0.3 mg/dL increase within 48 h	>0.5 and ≤ 1 mL/kg/h over 24 h
2	2.0–2.9 times baseline	<0.5 mL/kg/h for ≥12 h	2.0–2.9 times baseline*	>0.3 and ≤0.5 mL/kg/h over 24 h
3	≥3.0 times baseline OR Increase in serum creatinine to ≥4.0 mg/dL OR Initiation of renal replacement therapy OR Decrease in eGFR to <35 mL/min per 1.73 m ²	<0.3 mL/kg/h for ≥24 h OR Anuria for ≥12 h	≥3.0 times baseline* OR Increase in serum creatinine to ≥2.5 mg/dL OR Initiation of renal replacement therapy	≤0.3 mL/kg/h over 24 h

*Baseline serum creatinine is the lowest previous value.
Abbreviation: eGFR, estimated glomerular filtration rate.

was associated with poor prognosis^{20,21}. Urinary NGAL increases during urinary tract infection, sepsis, and CKD regardless of AKI²². A combination of two cell-cycle arrest markers urinary tissue inhibitor of metalloproteinase-2 and insulin-like growth factor-binding protein 7 demonstrated high sensitivity (90%) and low specificity (49%) in critically ill adults²³, and was approved by the United States Food and Drug Administration for use in adults. A few studies have shown that it has good diagnostic performance in pediatric, and neonatal AKI^{24,25}.

Despite the excellent performance of new biomarkers, they are not frequently used in clinical setting due to several concerns. First, they were validated for specific etiologies of AKI, such as pediatric patients undergoing cardiac surgery, and their efficacy cannot be reproduced in AKIs of different etiologies¹². Moreover, they are also elevated in other clinical conditions, such as inflammation¹². Large prospective studies investigating and validating the clinical utility of these biomarkers, therefore, is warranted.

Incidence

Two large, multinational epidemiological studies have been published in the pediatric and neonatal areas^{1,7}. The Assessment of Worldwide Acute Kidney Injury, Renal Angina, and Epidemiology (AWARE) study described the epidemiology of AKI in 4,683 pediatric patients (age range 3 months to 25 years) who were admitted to the PICU¹. It reported that AKI developed in 26.9% of patients and KDIGO stage 2 or 3 AKI in 11.6% within 7 days of PICU admission¹. The Assessment of Worldwide Acute Kidney injury Epidemiology in Neonates (AWAKEN) study described neonatal AKI in the neonatal ICU, and reported a prevalence similar to that of the AWARE study. The incidence of AKI was 29.9%, with 13.9% reaching AKI stage 1, 7.1% reaching AKI stage 2, and 8.9% reaching AKI stage 3⁷. The incidence outside the ICU varies across medical centers. To our knowledge, there have been no large-scale studies investigating AKI in non-critically ill children in Korea.

Outcomes

In the pediatric literature, AKI is consistently associated with poor outcomes, similar to adults. Many studies have demonstrated that AKI is an independent risk factor for prolonged stay in the PICU, longer duration of mechanical ventilation, and increased mortality among critically ill children^{1,26,27}. In addition, several observational studies have reported a high prevalence of CKD, hypertension, and proteinuria among AKI survivors²⁸⁻³⁰. Neonatal AKI is also a risk factor for mortality⁷. There have been no sufficiently large follow-up studies evaluating the relationship between neonatal AKI and long-term renal insufficiency. However, small observational studies have reported a high prevalence of CKD, and proteinuria in survivors of neonatal AKI^{29,31}.

Management

1. Risk factors and prevention

There are no effective medications for established AKI. Therefore, prevention and early detection are the mainstays of management. Monitoring high-risk patients and reducing additional risk factors can prevent the occurrence of AKI and improve outcomes. Prematurity, and chronic diseases such as CKD render the host susceptible to AKI, and events such as volume depletion, exposure to nephrotoxins, sepsis, major surgery, and critical illness lead to AKI⁴. The renal angina index was proposed to predict AKI in critically ill children on the basis of subtle kidney injury (changes in estimated creatinine clearance or fluid overload) and patient risk factors (ICU admission, stem cell transplantation, ventilation and inotropy)³².

Despite its importance, physicians often do not devote attention to the accumulation of risk factors and early decline of kidney function as they perform or attend to other important aspects of care³³. Recently, a system using electronic health records was implemented and helped physicians detect AKI early and mitigate the influence of risk factors³⁴. The system electronically flags high-risk patients to the medical team in near real-time. As such, the medical team does not miss high-risk patients and monitors them carefully. It has been shown to improve the rate of recovery from AKI³⁵. In the pediatric literature, this system was

developed to screen children who experienced multiple nephrotoxin exposures, which prompted clinicians to monitor more closely for the development of AKI. It demonstrated a positive effect in decreasing exposure to multiple nephrotoxins and, finally, AKI events³⁶.

2. Supportive care

Supportive care comprises optimization of volume status, blood pressure, avoidance of nephrotoxic agents, and nutritional support. It is important to maintain adequate renal perfusion through fluid and hemodynamic management³⁷. Volume status should be optimal (i.e., not excessive, not insufficient). The medical history and symptoms are important to evaluate volume status. Body weight, fluid intake, urine and stool output, and vital signs should be monitored daily, and lung sound and lower extremity edema should be checked^{38,39}. Serum chemistries and chest X-ray are also useful. Although volume depletion is a well-known risk factor, volume overload is associated with poor prognosis⁴⁰. Fluid accumulation in the acute phase, which is common in the PICU, is associated with high mortality in critically ill patients^{40,41}. The pediatric literature suggests that 10–20% fluid overload is a critical threshold at which outcomes are negatively impacted^{41,42}. Administration of optimal fluid amount is crucial. Patients with a normal intravascular volume should initially be limited to insensible losses (400 mL/m²/d) plus an amount of fluid equal to the urine output and extrarenal loss³⁸. Noticeable hypervolemic patients require further fluid restriction, omitting the replacement of insensible fluid losses, urine output, and extrarenal losses while considering adequate nutritional support³⁸. Diuretics therapy, especially loop diuretics, should be considered for hypervolemic patients. The pediatric literature shows that clinicians should consider initiating renal replacement therapy (RRT) at a fluid overload of >20%, while a fluid overload 10–20 % requires further evaluation⁴².

The proper type of fluid should be administered to prevent volume depletion. Crystalloids are the preferred solutions because data supporting the routine use of colloids for volume resuscitation are lacking⁴³. Colloid solution was not superior to crystalloid in terms of the prevention of AKI in children who underwent cardiac surgery⁴⁴. Among crystalloids, some studies have reported that a balanced

solution, such as Ringer's lactate solution, is superior to normal saline⁴⁵. Balanced solutions are defined as intravenous fluids having an electrolyte composition close to that of plasma. They have low chloride content of about 100–110 mmol/L, which is the most altered in normal saline as compared to plasma^{46,47}. Normal saline, which is hyperchloremic, can cause hyperchloremic metabolic acidosis, contributing to impaired recovery and poor clinical outcomes⁴⁸. Three recent large prospective studies have reported conflicting results^{49–51}; as such, debate regarding which solution is best among crystalloids continues.

Maintaining optimal blood pressure is crucial, although there is no currently definitive target value for pediatric AKI. For patients with hypotension, fluid resuscitation is initially considered if hypotension is due even partially to hypovolemia¹⁸. Norepinephrine is recommended if hypovolemia is not suspected⁵². High blood pressure in patients with AKI is mostly due to volume overload or disturbed renin-angiotensin axis²². Diuretics are attempted first unless patients exhibit signs of intravascular depletion²².

Nephrotoxin exposure is one of the most common causes of AKI in hospitalized children⁵³. Critically ill patients who are at high risk for AKI have the opportunity to be administered nephrotoxic medications. Amphotericin, aminoglycoside, vancomycin, nonsteroidal anti-inflammatory drugs, angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, calcineurin inhibitors, cisplatin, and methotrexate are well-known nephrotoxins¹⁸. Clinicians must balance the therapeutic benefits versus the risk for nephrotoxicity. If the use of nephrotoxic medications is inevitable, their dosage or dosing interval should be adjusted and monitored to reduce renal toxicity. Drug levels should be measured, if possible, and adjusted accordingly. If not possible, medications should be adjusted based on eGFR. Be aware that eGFR does not account for real-time renal function in AKI²². Contrast agent used for computed tomography and angiography are nephrotoxic. Contrast-induced AKI is common in children⁵⁴, and risk depends on the contrast agent dosage and type. Angiography requires more attention because it requires more contrast agent than computed tomography. High-osmolality contrast agent carries a higher risk for AKI, but have been replaced by low-osmolality contrast agents, which have a lower risk⁵⁵. Nevertheless, patients with a GFR <60 mL/

min/1.73m² or at high risk for AKI should be prepared before exposure²²). The key preventive measure is hydration. An isotonic crystalloid solution at 1 mL/kg/h should be administered intravenously for 12 hours before and after the procedure¹⁸. If contrast agent should be administered urgently, 3 mL/kg/h 1 h before and 6–9 mL/kg over 4–6 h after the procedure should be administered¹⁸. In addition, nephrotoxins should be discontinued for >24 h before the procedure¹⁸. The use of N-Acetylcysteine (NAC) will be discussed below.

Nutritional support is important for improving outcomes in children with AKI. For critically ill children with AKI, nutritional requirements should be individualized and assessed frequently⁵⁶. In general, 120–130% of basal calories needs, and 2–3 g/kg/day of protein should be provided⁵⁶, and hyperglycemia should be avoided⁵⁷.

3. Pharmacological treatment

Many researchers have investigated candidate medications to prevent and treat AKI. Anti-inflammatory, anti-oxidative and antiapoptotic interventions are representative examples; however, their yield remains insignificant. Although a few medications have demonstrated effectiveness in certain situations, most have yielded negative or conflicting results. Medications advocated in the past were not proven to be effective. The KDIGO guideline recommend NAC to prevent contrast-induced AKI in high-risk patients⁴. However, two large, well-designed studies reported no benefit of NAC in reducing the incidence of contrast-induced AKI^{58,59}. Diuretics are not recommended for the prevention of AKI because their use does not alter outcomes in those with established AKI⁶⁰. Diuretics should be used only to control fluid overload. Low-dose dopamine and fenoldopam did not have a positive effect on protection against AKI^{61,62}. Perioperative statins did not reduce the incidence of AKI among patients undergoing cardiac surgery⁶³. Even now, candidate medicines, such as recombinant alkaline phosphatase, costimulatory molecule CD28 receptor antagonist, p53 small interfering RNAs, and mesenchymal stem cells, are under consideration¹⁸. To establish the basis for routine clinical use of new medications, additional multicenter, high-quality trials are warranted.

4. RRT

RRT is required when conservative care fails. Indications for RRT include fluid overload (severe hypertension or pulmonary edema), severe hyperkalemia, metabolic acidosis, and severe uremia⁶⁴. The optimal timing of RRT initiation remains controversial. A recent meta-analysis of randomized controlled trials revealed that early initiation of RRT does not reduce mortality when compared with standard or late initiation in adults⁶⁵. Although one investigation involving pediatric patients demonstrated that early initiation improved mortality in cardiac surgery patients⁶⁶, it was small and retrospective in design; thus, more studies are needed. RRT modalities for AKI have begun with peritoneal dialysis and hemodialysis, and are now expanding to continuous renal replacement therapy (CRRT). We believe that all of these modalities represent viable management options in pediatric AKI⁶⁷. There is no definite evidence that one dialysis modality is superior to another in terms of outcomes in AKI^{68,69}. The modality should be determined according to the patient characteristics, institutional resources, and expertise⁶⁹. Each modality has advantages and disadvantages. Peritoneal dialysis can be used in small children such as preterm infants, but removes fluid and waste slowly and unpredictably⁶⁷. Hemodialysis removes toxins rapidly, but is dangerous for small children and hemodynamically unstable patients⁶⁷; additionally, it requires expertise. Currently, CRRT becomes the preferred modality in developed countries as improvement of equipment⁶⁹. CRRT facilitates hemodynamic stability and make it possible to use them in small children, such as neonates. New devices, such as the Cardio-Renal Pediatric Dialysis Emergency Machine, and Newcastle Infant Dialysis and Ultrafiltration System, have provided promising results for the treatment of neonates requiring RRT^{70,71}. However, the device should be operated in the PICU, so that children must endure anxiety about being separated from their parents during treatment.

Conclusion

AKI has attracted attention because of its high prevalence and association with poor outcomes. Many studies have been conducted to identify new biomarkers and effective

pharmacological interventions, and significant improvements have been achieved. Unfortunately, however, not many have been applied in actual clinical practice. Considering the gap between research achievements in adults and children, pediatric nephrologists have much to contribute to developments in understanding and treating AKI.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

ORCID

Myung Hyun Cho <https://orcid.org/0000-0002-3237-3173>.

Conflicts of interest

No potential conflict of interest relevant to this article was reported.

References

- Kaddourah A, Basu RK, Bagshaw SM, Goldstein SL. Epidemiology of Acute Kidney Injury in Critically Ill Children and Young Adults. *N Engl J Med* 2017;376:11-20.
- Madsen NL, Goldstein SL, Froslev T, Christiansen CF, Olsen M. Cardiac surgery in patients with congenital heart disease is associated with acute kidney injury and the risk of chronic kidney disease. *Kidney Int* 2017;92:751-6.
- Benisty K, Morgan C, Hessey E, Huynh L, Joffe AR, Garros D, et al. Kidney and blood pressure abnormalities 6 years after acute kidney injury in critically ill children: a prospective cohort study. *Pediatr Res* 2020.
- Kellum JA, Lameire N. Diagnosis, evaluation, and management of acute kidney injury: a KDIGO summary (Part 1). *Crit Care* 2013; 17:204.
- Zappitelli M, Parikh CR, Akcan-Arikan A, Washburn KK, Moffett BS, Goldstein SL. Ascertainment and epidemiology of acute kidney injury varies with definition interpretation. *Clin J Am Soc Nephrol* 2008;3:948-54.
- Charlton JR, Boohaker L, Askenazi D, Brophy PD, D'Angio C, Fuloria M, et al. Incidence and Risk Factors of Early Onset Neonatal AKI. *Clin J Am Soc Nephrol* 2019;14:184-95.
- Jetton JG, Boohaker LJ, Sethi SK, Wazir S, Rohatgi S, Soranno DE, et al. Incidence and outcomes of neonatal acute kidney injury (AWAKEN): a multicentre, multinational, observational cohort study. *Lancet Child Adolesc Health* 2017;1:184-94.
- Siegel SR, Oh W. Renal function as a marker of human fetal maturation. *Acta Paediatr Scand* 1976;65:481-5.
- Jetton JG, Askenazi DJ. Update on acute kidney injury in the neonate. *Curr Opin Pediatr* 2012;24:191-6.
- Bennett MR, Nehus E, Haffner C, Ma Q, Devarajan P. Pediatric reference ranges for acute kidney injury biomarkers. *Pediatr Nephrol* 2015;30:677-85.
- Devarajan P. Biomarkers for the early detection of acute kidney injury. *Curr Opin Pediatr* 2011;23:194-200.
- Beker BM, Corleto MG, Fieiras C, Musso CG. Novel acute kidney injury biomarkers: their characteristics, utility and concerns. *Int Urol Nephrol* 2018;50:705-13.
- Grubb A, Nyman U, Bjork J, Lindstrom V, Rippe B, Sterner G, et al. Simple cystatin C-based prediction equations for glomerular filtration rate compared with the modification of diet in renal disease prediction equation for adults and the Schwartz and the Counahan-Barratt prediction equations for children. *Clin Chem* 2005;51:1420-31.
- Hassinger AB, Backer CL, Lane JC, Haymond S, Wang D, Wald EL. Predictive power of serum cystatin C to detect acute kidney injury and pediatric-modified RIFLE class in children undergoing cardiac surgery. *Pediatr Crit Care Med* 2012;13:435-40.
- Lau L, Al-Ismailli Z, Harel-Sterling M, Pizzi M, Caldwell JS, Piccioni M, et al. Serum cystatin C for acute kidney injury evaluation in children treated with aminoglycosides. *Pediatr Nephrol* 2017;32: 163-71.
- El-Gammacy TM, Shinkar DM, Mohamed NR, Al-Halag AR. Serum cystatin C as an early predictor of acute kidney injury in preterm neonates with respiratory distress syndrome. *Scand J Clin Lab Invest* 2018;78:352-7.
- Benzer M, Alpay H, Baykan O, Erdem A, Demir IH. Serum NGAL, cystatin C and urinary NAG measurements for early diagnosis of contrast-induced nephropathy in children. *Ren Fail* 2016;38:27-34.
- Cho MH, Kang HG. Acute kidney injury and continuous renal replacement therapy in children; what pediatricians need to know. *Korean J Pediatr* 2018;61:339-47.
- Krawczeski CD, Goldstein SL, Woo JG, Wang Y, Piyaphanee N, Ma Q, et al. Temporal relationship and predictive value of urinary acute kidney injury biomarkers after pediatric cardiopulmonary bypass. *J Am Coll Cardiol* 2011;58:2301-9.
- Yoneyama F, Okamura T, Takigiku K, Yasukouchi S. Novel Urinary Biomarkers for Acute Kidney Injury and Prediction of Clinical Outcomes After Pediatric Cardiac Surgery. *Pediatr Cardiol* 2019.

21. Bennett M, Dent CL, Ma Q, Dastrala S, Grenier F, Workman R, et al. Urine NGAL predicts severity of acute kidney injury after cardiac surgery: a prospective study. *Clin J Am Soc Nephrol* 2008;3: 665-73.
22. Roy JP, Devarajan P. Acute Kidney Injury: Diagnosis and Management. *Indian J Pediatr* 2019.
23. Hall PS, Mitchell ED, Smith AF, Cairns DA, Messenger M, Hutchinson M, et al. The future for diagnostic tests of acute kidney injury in critical care: evidence synthesis, care pathway analysis and research prioritisation. *Health Technol Assess* 2018;22:1-274.
24. Westhoff JH, Tonshoff B, Waldherr S, Poschl J, Teufel U, Westhoff TH, et al. Urinary Tissue Inhibitor of Metalloproteinase-2 (TIMP-2) * Insulin-Like Growth Factor-Binding Protein 7 (IGFBP7) Predicts Adverse Outcome in Pediatric Acute Kidney Injury. *PLoS One* 2015;10:e0143628.
25. Chen J, Sun Y, Wang S, Dai X, Huang H, Bai Z, et al. The effectiveness of urinary TIMP-2 and IGFBP-7 in predicting acute kidney injury in critically ill neonates. *Pediatr Res* 2019.
26. Selewski DT, Cornell TT, Heung M, Troost JP, Ehrmann BJ, Lombel RM, et al. Validation of the KDIGO acute kidney injury criteria in a pediatric critical care population. *Intensive Care Med* 2014;40: 1481-8.
27. Alkandari O, Eddington KA, Hyder A, Gauvin F, Ducruet T, Gottesman R, et al. Acute kidney injury is an independent risk factor for pediatric intensive care unit mortality, longer length of stay and prolonged mechanical ventilation in critically ill children: a two-center retrospective cohort study. *Crit Care* 2011;15:R146.
28. Askenazi DJ, Feig DI, Graham NM, Hui-Stickle S, Goldstein SL. 3-5 year longitudinal follow-up of pediatric patients after acute renal failure. *Kidney Int* 2006;69:184-9.
29. Mammen C, Al Abbas A, Skippen P, Nadel H, Levine D, Collet JP, et al. Long-term risk of CKD in children surviving episodes of acute kidney injury in the intensive care unit: a prospective cohort study. *Am J Kidney Dis* 2012;59:523-30.
30. Menon S, Kirkendall ES, Nguyen H, Goldstein SL. Acute kidney injury associated with high nephrotoxic medication exposure leads to chronic kidney disease after 6 months. *J Pediatr* 2014;165:522-7.e2.
31. Polito C, Papale MR, La Manna A. Long-term prognosis of acute renal failure in the full-term neonate. *Clin Pediatr (Phila)* 1998;37: 381-5.
32. Basu RK, Zappitelli M, Brunner L, Wang Y, Wong HR, Chawla LS, et al. Derivation and validation of the renal angina index to improve the prediction of acute kidney injury in critically ill children. *Kidney Int* 2014;85:659-67.
33. Kumar Sethi S, Agrawal G, Wazir S, Rohatgi S, Iyengar A, Chakraborty R, et al. Neonatal Acute Kidney Injury: A Survey of Perceptions and Management Strategies Amongst Pediatricians and Neonatologists. *Front Pediatr* 2019;7:553.
34. Sutherland SM, Chawla LS, Kane-Gill SL, Hsu RK, Kramer AA, Goldstein SL, et al. Utilizing electronic health records to predict acute kidney injury risk and outcomes: workgroup statements from the 15(th) ADQI Consensus Conference. *Can J Kidney Health Dis* 2016;3:11.
35. Park S, Baek SH, Ahn S, Lee KH, Hwang H, Ryu J, et al. Impact of Electronic Acute Kidney Injury (AKI) Alerts With Automated Nephrologist Consultation on Detection and Severity of AKI: A Quality Improvement Study. *Am J Kidney Dis* 2018;71:9-19.
36. Goldstein SL, Kirkendall E, Nguyen H, Schaffzin JK, Bucuvalas J, Bracke T, et al. Electronic health record identification of nephrotoxin exposure and associated acute kidney injury. *Pediatrics* 2013;132:e756-67.
37. Chen H, Busse LW. Novel Therapies for Acute Kidney Injury. *Kidney Int Rep* 2017;2:785-99.
38. Devarajan P. Acute Kidney Injury. In: Kliegman RM, ST Geme III JW, Blum NJ, Shah SS, Tasker RC, Wilson KM, et al., editors. *Nelson Textbook of Pediatrics*. 21 ed. Philadelphia: Elsevier, 2019:2769-74.
39. Claire-Del Granado R, Mehta RL. Fluid overload in the ICU: evaluation and management. *BMC Nephrol* 2016;17:109.
40. Wang N, Jiang L, Zhu B, Wen Y, Xi XM. Fluid balance and mortality in critically ill patients with acute kidney injury: a multicenter prospective epidemiological study. *Crit Care* 2015;19:371.
41. Sutherland SM, Zappitelli M, Alexander SR, Chua AN, Brophy PD, Bunchman TE, et al. Fluid overload and mortality in children receiving continuous renal replacement therapy: the prospective pediatric continuous renal replacement therapy registry. *Am J Kidney Dis* 2010;55:316-25.
42. Selewski DT, Goldstein SL. The role of fluid overload in the prediction of outcome in acute kidney injury. *Pediatr Nephrol* 2018;33: 13-24.
43. Moore PK, Hsu RK, Liu KD. Management of Acute Kidney Injury: Core Curriculum 2018. *Am J Kidney Dis* 2018;72:136-48.
44. Oh HW, Lee JH, Kim HC, Kim EH, Song IK, Kim HS, et al. The effect of 6% hydroxyethyl starch (130/0.4) on acute kidney injury in paediatric cardiac surgery: a prospective, randomised trial. *Anaesthesia* 2018;73:205-15.
45. Krajewski ML, Raghunathan K, Paluszkiwicz SM, Schermer CR, Shaw AD. Meta-analysis of high- versus low-chloride content in perioperative and critical care fluid resuscitation. *Br J Surg* 2015; 102:24-36.
46. Hoorn EJ. Intravenous fluids: balancing solutions. *J Nephrol* 2017; 30:485-92.
47. Langer T, Santini A, Scotti E, Van Regenmortel N, Malbrain ML, Caironi P. Intravenous balanced solutions: from physiology to clinical evidence. *Anaesthesiol Intensive Ther* 2015;47 Spec No: 78-88.
48. Chowdhury AH, Cox EF, Francis ST, Lobo DN. A randomized, controlled, double-blind crossover study on the effects of 2-L infusions of 0.9% saline and plasma-lyte(R) 148 on renal blood flow velocity and renal cortical tissue perfusion in healthy volunteers. *Ann Surg* 2012;256:18-24.
49. Friederich A, Martin N, Swanson MB, Faine BA, Mohr NM. Normal Saline Solution and Lactated Ringer's Solution Have a Similar Ef-

- fect on Quality of Recovery: A Randomized Controlled Trial. *Ann Emerg Med* 2019;73:160-9.
50. Semler MW, Self WH, Wanderer JP, Ehrenfeld JM, Wang L, Byrne DW, et al. Balanced Crystalloids versus Saline in Critically Ill Adults. *N Engl J Med* 2018;378:829-39.
 51. Self WH, Semler MW, Wanderer JP, Wang L, Byrne DW, Collins SP, et al. Balanced Crystalloids versus Saline in Noncritically Ill Adults. *N Engl J Med* 2018;378:819-28.
 52. Joannidis M, Druml W, Forni LG, Groeneveld ABJ, Honore PM, Hoste E, et al. Prevention of acute kidney injury and protection of renal function in the intensive care unit: update 2017 : Expert opinion of the Working Group on Prevention, AKI section, European Society of Intensive Care Medicine. *Intensive Care Med* 2017;43:730-49.
 53. Hui-Stickle S, Brewer ED, Goldstein SL. Pediatric ARF epidemiology at a tertiary care center from 1999 to 2001. *Am J Kidney Dis* 2005;45:96-101.
 54. Cantais A, Hammouda Z, Mory O, Patural H, Stephan JL, Gulyaeva L, et al. Incidence of contrast-induced acute kidney injury in a pediatric setting: a cohort study. *Pediatr Nephrol* 2016;31:1355-62.
 55. Luk L, Steinman J, Newhouse JH. Intravenous Contrast-Induced Nephropathy-The Rise and Fall of a Threatening Idea. *Adv Chronic Kidney Dis* 2017;24:169-75.
 56. Sethi SK, Maxvold N, Bunchman T, Jha P, Kher V, Raina R. Nutritional management in the critically ill child with acute kidney injury: a review. *Pediatr Nephrol* 2017;32:589-601.
 57. Gordillo R, Ahluwalia T, Woroniecki R. Hyperglycemia and acute kidney injury in critically ill children. *Int J Nephrol Renovasc Dis* 2016;9:201-4.
 58. Weisbord SD, Gallagher M, Jneid H, Garcia S, Cass A, Thwin SS, et al. Outcomes after Angiography with Sodium Bicarbonate and Acetylcysteine. *N Engl J Med* 2018;378:603-14.
 59. ACT Investigators. Acetylcysteine for prevention of renal outcomes in patients undergoing coronary and peripheral vascular angiography: main results from the randomized Acetylcysteine for Contrast-induced nephropathy Trial (ACT). *Circulation* 2011;124:1250-9.
 60. Bagshaw SM, Delaney A, Haase M, Ghali WA, Bellomo R. Loop diuretics in the management of acute renal failure: a systematic review and meta-analysis. *Crit Care Resusc* 2007;9:60-8.
 61. Karthik S, Lisbon A. Low-dose dopamine in the intensive care unit. *Semin Dial* 2006;19:465-71.
 62. Bove T, Zangrillo A, Guarracino F, Alvaro G, Persi B, Maglioni E, et al. Effect of fenoldopam on use of renal replacement therapy among patients with acute kidney injury after cardiac surgery: a randomized clinical trial. *Jama* 2014;312:2244-53.
 63. Billings FTt, Hendricks PA, Schildcrout JS, Shi Y, Petracek MR, Byrne JG, et al. High-Dose Perioperative Atorvastatin and Acute Kidney Injury Following Cardiac Surgery: A Randomized Clinical Trial. *Jama* 2016;315:877-88.
 64. Sethi SK, Bunchman T, Raina R, Kher V. Unique considerations in renal replacement therapy in children: core curriculum 2014. *Am J Kidney Dis* 2014;63:329-45.
 65. Pasin L, Boraso S, Tiberio I. Early initiation of renal replacement therapy in critically ill patients: a meta-analysis of randomized clinical trials. *BMC Anesthesiol* 2019;19:62.
 66. Sanchez-de-Toledo J, Perez-Ortiz A, Gil L, Baust T, Lines-Palazon M, Perez-Hoyos S, et al. Early Initiation of Renal Replacement Therapy in Pediatric Heart Surgery Is Associated with Lower Mortality. *Pediatr Cardiol* 2016;37:623-8.
 67. Kwiatkowski DM, Sutherland SM. Acute kidney injury in pediatric patients. *Best Pract Res Clin Anaesthesiol* 2017;31:427-39.
 68. Nash DM, Przech S, Wald R, O'Reilly D. Systematic review and meta-analysis of renal replacement therapy modalities for acute kidney injury in the intensive care unit. *J Crit Care* 2017;41:138-44.
 69. de Galasso L, Picca S, Guzzo I. Dialysis modalities for the management of pediatric acute kidney injury. *Pediatr Nephrol* 2019.
 70. Ronco C, Garzotto F, Ricci Z. CA.R.PE.D.I.E.M. (Cardio-Renal Pediatric Dialysis Emergency Machine): evolution of continuous renal replacement therapies in infants. A personal journey. *Pediatr Nephrol* 2012;27:1203-11.
 71. Coulthard MG, Crosier J, Griffiths C, Smith J, Drinnan M, Whitaker M, et al. Haemodialysing babies weighing <8 kg with the Newcastle infant dialysis and ultrafiltration system (Nidus): comparison with peritoneal and conventional haemodialysis. *Pediatr Nephrol* 2014;29:1873-81.